



Общероссийский математический портал

N. P. Bondarenko, An inverse problem for the matrix quadratic pencil on a finite interval, *Eurasian Math. J.*, 2013, том 4, номер 3, 20–31

Использование Общероссийского математического портала Math-Net.Ru подразумевает, что вы прочитали и согласны с пользовательским соглашением
<http://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 18.97.14.83

23 января 2025 г., 04:01:30



AN INVERSE PROBLEM FOR THE MATRIX QUADRATIC PENCIL
ON A FINITE INTERVAL

N.P. Bondarenko

Communicated by Ya.T. Sultanaev

Key words: matrix quadratic differential pencils, Weyl matrix, inverse spectral problems, method of spectral mappings.

AMS Mathematics Subject Classification: 34A55, 34B07, 34B24, 34L40, 47E05.

Abstract. We consider a quadratic matrix boundary value problem with equations and boundary conditions dependent on a spectral parameter. We study an inverse problem that consists in recovering the differential pencil by the so-called Weyl matrix. We obtain asymptotic formulas for the solutions of the considered matrix equation. Using the ideas of the method of spectral mappings, we prove the uniqueness theorem for this inverse problem.

1 Introduction and main results

In this paper, we consider the boundary value problem $L = L(\ell, U, V)$ for the equation

$$\ell Y := Y'' + (\rho^2 \cdot I + 2i\rho Q_1(x) + Q_0(x))Y = 0, \quad x \in (0, \pi), \quad (1.1)$$

with the boundary conditions

$$\begin{aligned} U(Y) &:= Y'(0) + (i\rho h_1 + h_0)Y(0) = 0, \\ V(Y) &:= Y'(\pi) + (i\rho H_1 + H_0)Y(\pi) = 0. \end{aligned} \quad (1.2)$$

Here $Y(x) = [y_k(x)]_{k=\overline{1,m}}$ is a column vector, ρ is the spectral parameter, I is the $m \times m$ unit matrix, $Q_s(x) = [Q_{s,jk}(x)]_{j,k=\overline{1,m}}$ are $m \times m$ matrices with entries $Q_{s,jk}(x) \in W_1^s[0, \pi]$, $s = 0, 1$ $h_s = [h_{s,jk}]_{j,k=\overline{1,m}}$, $H_s = [H_{s,jk}]_{j,k=\overline{1,m}}$, where $h_{s,jk}$, $H_{s,jk}$ are complex numbers.

We assume that $\det(I \pm h_1) \neq 0$ and $\det(I \pm H_1) \neq 0$. This condition excludes problems of Regge type (see [23]) from consideration, as they require a separate investigation.

Differential equations with nonlinear dependence on the spectral parameter, or with so-called “energy-dependent” coefficients, frequently appear in mathematics and applications (see [15, 14, 16, 18, 21, 24] and references therein). In particular, inverse problems for such equations arise in investigation of mathematical models describing collisions of relativistic spinless particles [13] or proper vibrations of a string with viscous drag [22].

In this paper, we investigate the inverse problem for the pencil L , which consists in recovering coefficients of the boundary value problem (1.1), (1.2) by its spectral characteristics. In the scalar case ($m = 1$) inverse problems for quadratic pencils were studied in works [10, 25, 6, 7, 11, 20, 1, 2].

In the particular case when $Q_1(x) \equiv 0$, equation (1.1) becomes the matrix Sturm–Liouville equation. In recent years, significant progress has been made in the inverse problems theory for this equation. Constructive algorithms for solution of inverse problems were suggested, and characterization of spectral data was given (see [27, 8, 19, 3, 4]).

Now we plan to use ideas, developed for the matrix Sturm–Liouville equation, for problems with nonlinear dependence on the spectral parameter. In the present paper, we consider a general situation, without any conditions of selfadjointness on the coefficients and with the spectral parameter in the boundary conditions. We study the inverse problem for the pencil L by so-called Weyl matrix and prove the uniqueness theorem for the solution of this problem.

For our investigation, we develop ideas of the method of spectral mappings [9, 26]. This method also can be used to obtain a constructive procedure and numerical methods for the solution of this inverse problem. Numerical algorithms based on the method of spectral mappings are considered in the paper [12]. However, the questions of constructive solution of the inverse problem for the matrix pencil require separate investigation.

One of the main difficulties in problems for differential pencils is related to asymptotic behavior of solutions. The main terms of asymptotic representations depend on the coefficients of the pencil Q_1, h_1, H_1 . Derivation of asymptotic formulas is nontrivial even in the scalar case, and in the matrix case it is more complicated, because of additional difficulties related to noncommutativity of matrix multiplication. We obtain the asymptotics for the fundamental system of solutions for equation (1.1). They are required to prove uniqueness, but they can also be considered as a separate result.

Now we proceed to the formulation of the main result. Let $\Phi(x, \rho) = [\Phi_{jk}(x, \rho)]_{j,k=\overline{1,m}}$ be the matrix solution of equation (1.1) satisfying the conditions $U(\Phi) = I, V(\Phi) = 0$. We call $\Phi(x, \rho)$ the *Weyl solution* for L . Put $M(\rho) := \Phi(0, \rho)$. The matrix $M(\rho) = [M_{jk}(\rho)]_{j,k=\overline{1,m}}$ is called the *Weyl matrix* for L . The notion of the Weyl matrix is a generalization of the notion of the Weyl function (m -function) for the scalar case (see [17, 9]) and the notion of the Weyl matrix for the matrix Sturm–Liouville operator (see [3, 4]).

The inverse problem is stated as follows.

Inverse Problem 1. *Given a Weyl matrix $M(\rho)$, find the coefficients of the pencil L .*

Remark. One can prove in a standard way that the boundary problem L has a countable set of eigenvalues $\{\rho_n\}$. The entries of $M(\rho)$ are meromorphic in ρ and their poles coincide with $\{\rho_n\}$. As in the scalar case (see [6]), the following representation is valid

$$M(\rho) = \sum_n \sum_{\nu=1}^{m_n} \frac{M_{n\nu}}{(\rho - \rho_n)^\nu}, \quad (1.3)$$

where m_n are multiplicities of the corresponding eigenvalues ρ_n , and $M_{n\nu}$ are some matrix coefficients.

Following [6, 7], we call the collection $\{\rho_n, M_{n\nu}\}$ the *spectral data* of the pencil L . By virtue of (1.3), the spectral data determine the Weyl matrix uniquely. Therefore, Inverse Problem 1 is equivalent to

Inverse Problem 2. Given spectral data $\{\rho_n, M_{n\nu}\}$, find the coefficients of the pencil L .

In this paper, we restrict ourselves to Inverse Problem 1.

Along with L we consider a pencil \tilde{L} of the same form but with other coefficients $\tilde{Q}_s(x)$, \tilde{h}_s , \tilde{H}_s . We agree that if a symbol γ denotes an object related to L then $\tilde{\gamma}$ denotes the corresponding object related to \tilde{L} .

We now state the uniqueness theorem for Inverse Problem 1.

Theorem 1.1. *If $M(\rho) = \tilde{M}(\rho)$, then $L = \tilde{L}$. Hence the Weyl matrix determines the coefficients of the pencil (1.1), (1.2) uniquely.*

In order to prove Theorem 1.1, we need asymptotics for the solutions to equation (1.1). They are obtained in Section 2. In Section 3, we provide the proof of the uniqueness theorem.

2 Asymptotic behaviour of the solutions

Let $C(x, \rho) = [C_{jk}(x, \rho)]_{j,k=\overline{1,m}}$ and $S(x, \rho) = [S_{jk}(x, \rho)]_{j,k=\overline{1,m}}$ be the matrix solutions of equation (1.1) under the initial conditions

$$C(0, \rho) = S'(0, \rho) = I, \quad C'(0, \rho) = S(0, \rho) = 0.$$

The main goal of this section is to obtain the asymptotics of $S(x, \rho)$ and $C(x, \rho)$ as $|\rho| \rightarrow \infty$.

Let the matrix functions $P_+(x)$, $P_-(x)$, $P_+^*(x)$ and $P_-^*(x)$ be the solutions of the Cauchy problems

$$\begin{aligned} P'_\pm(x) &= \pm Q_1(x)P_\pm(x), & P_\pm(0) &= I, \\ P'^*_\pm(x) &= \pm P^*_\pm(x)Q_1(x), & P^*_\pm(0) &= I. \end{aligned} \tag{2.1}$$

Remark. In the scalar case ($m = 1$), one has

$$P_\pm(x) = P^*_\pm(x) = \exp \left\{ \pm \int_0^x Q_1(t) dt \right\}.$$

Lemma 2.1. *The following relations hold*

$$P_+(x)P_-^*(x) = P_-^*(x)P_+(x) = I, \quad P_-(x)P_+^*(x) = P_+^*(x)P_-(x) = I,$$

for all $x \in [0, \pi]$.

Proof. Using (2.1), we get

$$\begin{aligned} (P_-^*(x)P_+(x))' &= (P_-^*(x))'P_+(x) + P_-^*(x)P_+'(x) \\ &= -P_-^*(x)Q_1(x)P_+(x) + P_-^*(x)Q_1(x)P_+(x) = 0. \end{aligned}$$

Hence $P_-^*(x)P_+(x)$ does not depend on x , so $P_-^*(x)P_+(x) = P_-^*(0)P_+(0) = I$. The other relations can be proved similarly. \square

Theorem 2.1. *For $x \in [0, \pi]$, $|\rho| \rightarrow \infty$, $\nu = 0, 1$, the following relations hold*

$$C^{(\nu)}(x, \rho) = \frac{(i\rho)^\nu}{2} \exp(i\rho x)P_-(x) + \frac{(-i\rho)^\nu}{2} \exp(-i\rho x)P_+(x) + O(\rho^{\nu-1} \exp(|\tau|x)), \quad (2.2)$$

$$S^{(\nu)}(x, \rho) = \frac{(i\rho)^{\nu-1}}{2} \exp(i\rho x)P_-(x) + \frac{(-i\rho)^{\nu-1}}{2} \exp(-i\rho x)P_+(x) + O(\rho^{\nu-2} \exp(|\tau|x)),$$

where $\tau := \text{Im } \rho$.

In order to prove the theorem, we develop the ideas of [5] with necessary modifications caused by the matrix case.

Proof. 1. First we derive Volterra integral equations for $S(x, \rho)$ and $C(x, \rho)$.

One can easily check that the matrix functions

$$\begin{aligned} C_0(x, \rho) &= \frac{1}{2} \exp(i\rho x)P_-(x) + \frac{1}{2} \exp(-i\rho x)P_+(x), \\ S_0(x, \rho) &= \frac{1}{2i\rho} \exp(i\rho x)P_-(x) - \frac{1}{2i\rho} \exp(-i\rho x)P_+(x) \end{aligned}$$

form a fundamental system of solutions for the differential equation

$$Y'' + Q_1'(x)(i\rho \cdot I - Q_1(x))^{-1}Y' + (\rho^2 \cdot I + 2i\rho Q_1(x) - Q_1^2(x))Y = 0.$$

Rewrite equation (1.1) in the form

$$Y'' + Q_1'(x)(i\rho \cdot I - Q_1(x))^{-1}Y' + (\rho^2 \cdot I + 2i\rho Q_1(x) - Q_1^2(x))Y = F(x, \rho, Y), \quad (2.3)$$

$$F(x, \rho, Y) := Q_1'(x)(i\rho \cdot I - Q_1(x))^{-1}Y' - (Q_1^2(x) + Q_0(x))Y. \quad (2.4)$$

Apply the method of variation of parameters to this equation. Every solution of (2.3) can be represented in the form

$$Y(x, \rho) = C_0(x, \rho)A(x, \rho) + S_0(x, \rho)B(x, \rho),$$

where coefficient matrices $A(x, \rho)$ and $B(x, \rho)$ can be found from the system

$$\begin{bmatrix} C_0(x, \rho) & S_0(x, \rho) \\ C_0'(x, \rho) & S_0'(x, \rho) \end{bmatrix} \cdot \begin{bmatrix} A'(x, \rho) \\ B'(x, \rho) \end{bmatrix} = \begin{bmatrix} 0 \\ F(x, \rho, Y) \end{bmatrix} \quad (2.5)$$

and the initial conditions

$$A(0, \rho) = Y(0, \rho), \quad B(0, \rho) = (I - Q_1(0)/(i\rho))^{-1}Y'(0, \rho). \quad (2.6)$$

In order to solve system (2.5), we introduce the matrix functions

$$\begin{aligned} C_0^*(x, \rho) &= \frac{1}{2} \exp(i\rho x)P_-^*(x) + \frac{1}{2} \exp(-i\rho x)P_+^*(x), \\ S_0^*(x, \rho) &= \frac{1}{2i\rho} \exp(i\rho x)P_-^*(x) - \frac{1}{2i\rho} \exp(-i\rho x)P_+^*(x). \end{aligned}$$

Using Lemma 2.1, one can easily show that

$$\begin{bmatrix} C_0 & S_0 \\ C_0' & S_0' \end{bmatrix}^{-1} = \begin{bmatrix} S_0^{*'} & -S_0^* \\ -C_0^{*'} & C_0^* \end{bmatrix} \cdot \begin{bmatrix} i\rho(i\rho \cdot I - Q_1)^{-1} & 0 \\ 0 & i\rho(i\rho \cdot I - Q_1)^{-1} \end{bmatrix}.$$

Therefore

$$A'(x, \rho) = -i\rho S_0^*(x, \rho)(i\rho \cdot I - Q_1(x))^{-1}F(x, \rho, Y),$$

$$B'(x, \rho) = i\rho C_0^*(x, \rho)(i\rho \cdot I - Q_1(x))^{-1}F(x, \rho, Y).$$

and

$$\begin{aligned} Y(x, \rho) &= C_0(x, \rho)A(0, \rho) + S_0(x, \rho)B(0, \rho) + \\ &+ i\rho \int_0^x (S_0(x, \rho)C_0^*(t, \rho) - C_0(x, \rho)S_0^*(t, \rho))(i\rho \cdot I - Q_1(t))^{-1}F(t, \rho, Y) dt. \end{aligned} \quad (2.7)$$

Simple calculations show that

$$\begin{aligned} &S_0(x, \rho)C_0^*(t, \rho) - C_0(x, \rho)S_0^*(t, \rho) \\ &= \frac{1}{2i\rho} \{ \exp(i\rho(x-t))P_-(x)P_+^*(t) - \exp(-i\rho(x-t))P_+(x)P_-^*(t) \}. \end{aligned}$$

Substituting $Y = C(x, \rho)$ and $Y = S(x, \rho)$ into (2.6) and (2.7), we arrive at the following integral equations

$$\begin{aligned} C(x, \rho) &= C_0(x, \rho) + \frac{1}{2} \int_0^x \{ \exp(i\rho(x-t))P_-(x)P_+^*(t) - \\ &- \exp(-i\rho(x-t))P_+(x)P_-^*(t) \} (i\rho \cdot I - Q_1(t))^{-1}F(t, \rho, C) dt, \end{aligned} \quad (2.8)$$

$$\begin{aligned} S(x, \rho) &= S_0(x, \rho) \left(I - \frac{Q_1(0)}{i\rho} \right)^{-1} + \frac{1}{2} \int_0^x \{ \exp(i\rho(x-t))P_-(x)P_+^*(t) - \\ &- \exp(-i\rho(x-t))P_+(x)P_-^*(t) \} (i\rho \cdot I - Q_1(t))^{-1}F(t, \rho, S) dt, \end{aligned}$$

where $F(x, \rho, Y)$ is defined in (2.4).

2. Then we continue to work with $C(x, \rho)$. The function $S(x, \rho)$ can be treated similarly. Differentiating (2.8) with respect to x and using (2.1), we get

$$C'(x, \rho) = C'_0(x, \rho) + \frac{1}{2}(i\rho \cdot I - Q_1(x)) \int_0^x \{ \exp(i\rho(x-t))P_-(x)P_+^*(t) + \exp(-i\rho(x-t))P_+(x)P_-^*(t) \} (i\rho \cdot I - Q_1(t))^{-1} F(t, \rho, C) dt, \quad (2.9)$$

Denote

$$\mu_\nu(\rho) := \max_{x \in [0, \pi]} \|C^{(\nu)}(x, \rho) \exp(-|\tau|x)\|, \quad \nu = 0, 1.$$

We agree to denote by the same symbol K different positive constants not depending on x and ρ , and we mean by $\|\cdot\|$ the following matrix norm:

$$\|A\| = \max_{j=1, m} \sum_{k=1}^m |a_{jk}|,$$

$$A = [a_{jk}]_{j, k=1, \overline{m}}.$$

Since

$$\|F(t, \rho, C)\| \leq K \|Q'_1(t)\| \frac{\mu_1(\rho)}{|\rho|} + \|Q_1^2(t) + Q_0(t)\| \mu_0(\rho),$$

we get from (2.8) and (2.9)

$$\mu_0(\rho) \leq K \left(1 + \frac{\mu_0(\rho)}{|\rho|} + \frac{\mu_1(\rho)}{|\rho|^2} \right), \quad \mu_1(\rho) \leq K \left(|\rho| + \mu_0(\rho) + \frac{\mu_1(\rho)}{|\rho|} \right),$$

whence we obtain $\mu_\nu(\rho) \leq K|\rho|^\nu$ or

$$C^{(\nu)}(x, \rho) = O(|\rho|^\nu \exp(|\tau|x)).$$

Substituting this into (2.8) and (2.9), we arrive at (2.2). \square

3 Proof of the uniqueness theorem

Let $\varphi(x, \rho) = [\varphi_{jk}(x, \rho)]_{j, k=1, \overline{m}}$ and $\psi(x, \rho) = [\psi_{jk}(x, \rho)]_{j, k=1, \overline{m}}$ be the matrix solutions of equation (1.1) under the initial conditions

$$\varphi(0, \rho) = \psi(\pi, \rho) = I, \quad U(\varphi) = V(\psi) = 0.$$

It is easy to check that

$$\Phi(x, \rho) = S(x, \rho) + \varphi(x, \rho)M(\rho), \quad (3.1)$$

$$\Phi(x, \rho) = \psi(x, \rho)(U(\psi))^{-1}. \quad (3.2)$$

We expand $\varphi(x, \rho)$ by the fundamental system of solutions $C(x, \rho)$ and $S(x, \rho)$

$$\varphi(x, \rho) = C(x, \rho) - S(x, \rho)(i\rho h_1 + h_0)$$

and, using Theorem 2.1, we get

$$\begin{aligned} \varphi^{(\nu)}(x, \rho) &= \frac{(i\rho)^\nu}{2} \exp(i\rho x) P_-(x)(I - h_1) + \frac{(-i\rho)^\nu}{2} \exp(-i\rho x) P_+(x)(I + h_1) \\ &\quad + O(\rho^{\nu-1} \exp(|\tau|x)), \quad x \in [0, \pi], \nu = 0, 1, |\rho| \rightarrow \infty. \end{aligned}$$

In order to obtain the asymptotics for $\psi(x, \rho)$, we can apply the substitution $x \rightarrow \pi - x$, $h_1 \rightarrow -H_1$, $h_0 \rightarrow -H_0$. Then we need the analogs of $P_\pm(x)$ and $P_\pm^*(x)$, which we denote by $P_\pm^\bullet(x)$ and $P_\pm^{\bullet*}(x)$ and define as the solutions of the Cauchy problems

$$\begin{aligned} P_\pm^{\bullet'}(x) &= \mp Q_1(x) P_\pm^\bullet(x), \quad P_\pm^\bullet(\pi) = I, \\ P_\pm^{\bullet*'}(x) &= \mp P_\pm^{\bullet*}(x) Q_1(x), \quad P_\pm^{\bullet*}(\pi) = I. \end{aligned}$$

The following lemma establishes connections between these asymptotic coefficients.

Lemma 3.1. *For $x \in [0, \pi]$, the following relations hold*

$$\begin{aligned} P_+(x) P_-^\bullet(0) &= P_-^\bullet(x), \quad P_-(x) P_+^\bullet(0) = P_+^\bullet(x), \\ P_-^{\bullet*}(0) P_+^*(x) &= P_-^{\bullet*}(x), \quad P_+^{\bullet*}(0) P_-^*(x) = P_+^{\bullet*}(x). \end{aligned}$$

The proof, based on using the corresponding Cauchy problems, is trivial.

One can obtain the following asymptotic formula

$$\begin{aligned} \psi^{(\nu)}(x, \rho) &= \frac{(-i\rho)^\nu}{2} \exp(i\rho(\pi - x)) P_-^\bullet(x)(I + H_1) + \frac{(i\rho)^\nu}{2} \exp(-i\rho(\pi - x)) P_+^\bullet(x)(I - H_1) \\ &\quad + O(\rho^{\nu-1} \exp(|\tau|(\pi - x))), \quad x \in [0, \pi], \nu = 0, 1, |\rho| \rightarrow \infty, \end{aligned}$$

and then using (3.2), one can derive asymptotic formulas for $\Phi(x, \rho)$ and $M(\rho)$.

Hereafter the asymptotics in the angles $\Theta_\delta^\pm := \{\rho \in \mathbb{C} : \delta \leq \pm \arg \rho \leq \pi - \delta\}$, $0 < \delta < \pi$ will be required, so we formulate the following result.

Lemma 3.2. *Suppose $x \in (0, \pi)$, $\nu = 0, 1$, $|\rho| \rightarrow \infty$; then*

(i) *for $\rho \in \Theta_\delta^+$, we have*

$$\begin{aligned} \varphi^{(\nu)}(x, \rho) &= \frac{(-i\rho)^\nu}{2} \exp(-i\rho x) P_+(x)(I + h_1) + O(\rho^{\nu-1} \exp(|\tau|x)), \\ \psi^{(\nu)}(x, \rho) &= \frac{(i\rho)^\nu}{2} \exp(-i\rho(\pi - x)) P_+^\bullet(x)(I - H_1) + O(\rho^{\nu-1} \exp(|\tau|(\pi - x))), \\ \Phi^{(\nu)}(x, \rho) &= -(-i\rho)^{\nu-1} \exp(i\rho x) P_+^\bullet(x) (P_+^\bullet(0))^{-1} (I + h_1)^{-1} + O(\rho^{\nu-2} \exp(-|\tau|x)), \\ M(\rho) &= (i\rho)^{-1} (I + h_1)^{-1} + O(\rho^{-2}). \end{aligned}$$

(ii) *for $\rho \in \Theta_\delta^-$, we have*

$$\begin{aligned} \varphi^{(\nu)}(x, \rho) &= \frac{(i\rho)^\nu}{2} \exp(i\rho x) P_-(x)(I - h_1) + O(\rho^{\nu-1} \exp(|\tau|x)), \\ \psi^{(\nu)}(x, \rho) &= \frac{(-i\rho)^\nu}{2} \exp(i\rho(\pi - x)) P_-^\bullet(x)(I + H_1) + O(\rho^{\nu-1} \exp(|\tau|(\pi - x))), \\ \Phi^{(\nu)}(x, \rho) &= -(i\rho)^{\nu-1} \exp(-i\rho x) P_-^\bullet(x) (P_-^\bullet(0))^{-1} (I - h_1)^{-1} + O(\rho^{\nu-2} \exp(-|\tau|x)), \\ M(\rho) &= -(i\rho)^{-1} (I - h_1)^{-1} + O(\rho^{-2}). \end{aligned}$$

By definition, put

$$\begin{aligned}\ell^*Z &:= Z'' + Z(\rho^2 \cdot I + 2i\rho Q_1(x) + Q_0(x)), \\ U^*(Z) &:= Z'(0) + Z(0)(i\rho h_1 + h_0), \\ V^*(Z) &:= Z'(\pi) + Z(\pi)(i\rho H_1 + H_0), \\ \langle Z, Y \rangle &= Z'(x)Y(x) - Z(x)Y'(x),\end{aligned}$$

where $Z = [z_k]_{k=1, \overline{m}}^t$ is a row vector (t is the sign for the transposition). Then

$$\langle Z, Y \rangle|_{x=0} = U^*(Z)Y(0) - Z(0)U(Y), \quad \langle Z, Y \rangle|_{x=\pi} = V^*(Z)Y(\pi) - Z(\pi)V(Y). \quad (3.3)$$

If $Y(x, \rho)$ and $Z(x, \rho)$ satisfy the equations $\ell Y(x, \rho) = 0$ and $\ell^* Z(x, \rho) = 0$ respectively, then

$$\frac{d}{dx} \langle Z(x, \rho), Y(x, \rho) \rangle = 0. \quad (3.4)$$

Let $\varphi^*(x, \rho)$, $S^*(x, \rho)$, $\psi^*(x, \rho)$ and $\Phi^*(x, \rho)$ be the matrices, satisfying the equation $\ell^* Z = 0$ and the conditions

$$\varphi^*(0, \rho) = S^{*'}(0, \rho) = \psi^*(\pi, \rho) = U^*(\Phi^*) = I,$$

$U^*(\varphi^*) = S^*(0, \rho) = V^*(\psi^*) = V^*(\Phi^*) = 0$. Put $M^*(\rho) := \Phi^*(0, \rho)$. Then

$$\Phi^*(x, \rho) = S^*(x, \rho) + M^*(\rho)\varphi^*(x, \rho) = (U^*(\psi^*))^{-1}\psi^*(x, \rho). \quad (3.5)$$

According to (3.4), $\langle \Phi^*(x, \rho), \Phi(x, \rho) \rangle$ does not depend on x . Using (3.3), we get

$$\langle \Phi^*(x, \rho), \Phi(x, \rho) \rangle|_{x=0} = M(\rho) - M^*(\rho), \quad \langle \Phi^*(x, \rho), \Phi(x, \rho) \rangle|_{x=\pi} = 0.$$

Therefore, $M(\rho) \equiv M^*(\rho)$.

Using (3.3), one can easily show that

$$\begin{bmatrix} \Phi^{*'}(x, \rho) & -\Phi^*(x, \rho) \\ -\varphi^{*'}(x, \rho) & \varphi^*(x, \rho) \end{bmatrix} \begin{bmatrix} \varphi(x, \rho) & \Phi(x, \rho) \\ \varphi'(x, \rho) & \Phi'(x, \rho) \end{bmatrix} = \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix}.$$

Hence,

$$\begin{bmatrix} \varphi(x, \rho) & \Phi(x, \rho) \\ \varphi'(x, \rho) & \Phi'(x, \rho) \end{bmatrix}^{-1} = \begin{bmatrix} \Phi^{*'}(x, \rho) & -\Phi^*(x, \rho) \\ -\varphi^{*'}(x, \rho) & \varphi^*(x, \rho) \end{bmatrix}. \quad (3.6)$$

We also need the asymptotics for $\varphi^*(x, \rho)$ and $\Phi^*(x, \rho)$.

Lemma 3.3. *Suppose $x \in (0, \pi)$, $\nu = 0, 1$, $|\rho| \rightarrow \infty$; then*

(i) *for $\rho \in \Theta_\delta^+$, we have*

$$\varphi^{*(\nu)}(x, \rho) = \frac{(-i\rho)^\nu}{2} \exp(-i\rho x)(I + h_1)P_+^*(x) + O(\rho^{\nu-1} \exp(|\tau|x)),$$

$$\psi^{*(\nu)}(x, \rho) = \frac{(i\rho)^\nu}{2} \exp(-i\rho(\pi - x))(I - H_1)P_+^{*\bullet}(x) + O(\rho^{\nu-1} \exp(|\tau|(\pi - x))),$$

$$\Phi^{*(\nu)}(x, \rho) = -(-i\rho)^{\nu-1} \exp(i\rho x)(I + h_1)^{-1}(P_+^{\bullet*}(0))^{-1}P_+^{\bullet*}(x) + O(\rho^{\nu-2} \exp(-|\tau|x)).$$

(ii) for $\rho \in \Theta_\delta^-$, we have

$$\varphi^{*(\nu)}(x, \rho) = \frac{(i\rho)^\nu}{2} \exp(i\rho x)(I - h_1)P_-^*(x) + O(\rho^{\nu-1} \exp(|\tau|x)),$$

$$\psi^{*(\nu)}(x, \rho) = \frac{(-i\rho)^\nu}{2} \exp(i\rho(\pi - x))(I + H_1)P_-^{\bullet*}(x) + O(\rho^{\nu-1} \exp(|\tau|(\pi - x))),$$

$$\Phi^{*(\nu)}(x, \rho) = -(i\rho)^{\nu-1} \exp(-i\rho x)(I - h_1)^{-1}(P_-^{\bullet*}(0))^{-1}P_-^{\bullet*}(x) + O(\rho^{\nu-2} \exp(-|\tau|x)).$$

Proof of Theorem 1.1. Consider the problems L and \tilde{L} with the Weyl matrices $M(\lambda) \equiv \tilde{M}(\lambda)$. Note that according to asymptotics for $M(\lambda)$ of Lemma 3.2, we straightway get

$$h_1 = \tilde{h}_1. \quad (3.7)$$

Now we consider the block-matrix $\mathcal{P}(x, \rho) = [\mathcal{P}_{jk}(x, \rho)]_{j,k=1,2}$ defined by

$$\mathcal{P}(x, \rho) \begin{bmatrix} \tilde{\varphi}(x, \rho) & \tilde{\Phi}(x, \rho) \\ \tilde{\varphi}'(x, \rho) & \tilde{\Phi}'(x, \rho) \end{bmatrix} = \begin{bmatrix} \varphi(x, \rho) & \Phi(x, \rho) \\ \varphi'(x, \rho) & \Phi'(x, \rho) \end{bmatrix}. \quad (3.8)$$

Taking (3.6) into account, we calculate

$$\begin{aligned} \mathcal{P}_{j1}(x, \rho) &= \varphi^{(j-1)}(x, \rho)\tilde{\Phi}^{*'}(x, \rho) - \Phi^{(j-1)}(x, \rho)\tilde{\varphi}^{*'}(x, \rho), \\ \mathcal{P}_{j2}(x, \rho) &= \Phi^{(j-1)}(x, \rho)\tilde{\varphi}^*(x, \rho) - \varphi^{(j-1)}(x, \rho)\tilde{\Phi}^*(x, \rho). \end{aligned} \quad (3.9)$$

Applying Lemmas 3.2 and 3.3 and using (3.7), for $\rho \in \Theta_\delta^+$, $|\rho| \rightarrow \infty$, $x \in (0, \pi)$ we calculate

$$\begin{aligned} \mathcal{P}_{11}(x, \rho) &= \frac{1}{2}P_+(x)(\tilde{P}_+^{\bullet*}(0))^{-1}\tilde{P}_+^{\bullet*}(x) + \frac{1}{2}P_+^\bullet(x)(\tilde{P}_+^*(0))^{-1}\tilde{P}_+^*(x) + O(\rho^{-1}), \\ \mathcal{P}_{12}(x, \rho) &= \frac{1}{2i\rho}P_+^\bullet(x)(P_+^\bullet(0))^{-1}\tilde{P}_+^*(x) - \frac{1}{2i\rho}P_+(x)(\tilde{P}_+^{\bullet*}(0))^{-1}\tilde{P}_+^{\bullet*}(x) + O(\rho^{-2}). \end{aligned}$$

Similarly, for $\rho \in \Theta_\delta^-$, $|\rho| \rightarrow \infty$, $x \in (0, \pi)$

$$\begin{aligned} \mathcal{P}_{11}(x, \rho) &= \frac{1}{2}P_-(x)(\tilde{P}_-^{\bullet*}(0))^{-1}\tilde{P}_-^{\bullet*}(x) + \frac{1}{2}P_-^\bullet(x)(\tilde{P}_-^*(0))^{-1}\tilde{P}_-^*(x) + O(\rho^{-1}), \\ \mathcal{P}_{12}(x, \rho) &= -\frac{1}{2i\rho}P_-^\bullet(x)(P_-^\bullet(0))^{-1}\tilde{P}_-^*(x) + \frac{1}{2i\rho}P_-(x)(\tilde{P}_-^{\bullet*}(0))^{-1}\tilde{P}_-^{\bullet*}(x) + O(\rho^{-2}). \end{aligned}$$

Using Lemma 3.1, we obtain for $\rho \in \Theta_\delta^+ \cup \Theta_\delta^-$, $|\rho| \rightarrow \infty$, $x \in (0, \pi)$, that

$$\mathcal{P}_{11}(x, \rho) = \Omega(x) + O(\rho^{-1}), \quad \mathcal{P}_{12}(x, \rho) = \rho^{-1}\Lambda(x) + O(\rho^{-2}), \quad (3.10)$$

where

$$\Omega(x) := \frac{1}{2} \left(P_-(x)\tilde{P}_+^*(x) + P_+(x)\tilde{P}_-^*(x) \right),$$

$$\Lambda(x) := \frac{1}{2i} \left(P_-(x) \tilde{P}_+^*(x) - P_+(x) \tilde{P}_-^*(x) \right).$$

Substituting (3.1) and (3.5) into (3.9), we get

$$\mathcal{P}_{11} = \varphi \tilde{S}^{*'} - S \tilde{\varphi}^{*'} + \varphi (\tilde{M}^* - M) \tilde{\varphi}^*,$$

$$\mathcal{P}_{12} = S \tilde{\varphi}^* - \varphi \tilde{S}^* + \varphi (M - \tilde{M}^*) \tilde{\varphi}^*,$$

Note that for each fixed $x \in (0, \pi)$, the matrix functions φ , φ^* , S , S^* and they derivatives with respect to x are entire in ρ of order 1. Since $\tilde{M}^*(\rho) \equiv \tilde{M}(\rho) \equiv M(\rho)$, it follows for each fixed $x \in (0, \pi)$ that the entries of $\mathcal{P}_{11}(x, \rho)$ and $\mathcal{P}_{12}(x, \rho)$ are entire functions in ρ of order not greater than 1. Together with (3.10) this yields $\mathcal{P}_{11}(x, \rho) \equiv \Omega(x)$, $\mathcal{P}_{12}(x, \rho) \equiv 0$, $\Lambda(x) \equiv 0$.

Differentiating $\Omega(x)$ and using (2.1), we get

$$\Omega'(x) = i(\Lambda(x) \tilde{Q}_1(x) - Q_1(x) \Lambda(x)) = 0$$

for almost all x in $[0, \pi]$. Therefore, $\Omega(x) \equiv \Omega(0) \equiv I$, $x \in [0, \pi]$. Thus, $\mathcal{P}_{11}(x, \rho) = I$, $x \in (0, \pi)$. By virtue of (3.8) we have $\varphi(x, \rho) \equiv \tilde{\varphi}(x, \rho)$, $\Phi(x, \rho) \equiv \tilde{\Phi}(x, \rho)$ and consequently, $L = \tilde{L}$. \square

Acknowledgments

This research was supported in part by the German Academic Exchange Service (DAAD) and Grant 13-01-00134 of the Russian Foundation for Basic Research.

References

- [1] A.M. Akhtyamov, V.A. Sadovnichy, Ya.T. Sultanaev, *Inverse problem for an operator pencil with nonseparated boundary conditions*, Eurasian Math. J. 1 (2010), no. 2, 5–16.
- [2] A.M. Akhtyamov, V.A. Sadovnichy, Ya.T. Sultanaev, *Generalizations of Borg’s uniqueness theorem to the case of nonseparated boundary conditions*, Eurasian Math. J. 3 (2012), no. 4, 10–22.
- [3] N. Bondarenko, *Spectral analysis for the matrix Sturm-Liouville operator on a finite interval*, Tamkang J. Math. 42 (2011), no. 3, 305-327.
- [4] N.P. Bondarenko, *Necessary and sufficient conditions for the solvability of the inverse problem for the matrix Sturm-Liouville operator*, Funct. Anal. Appl. 46 (2012), no. 1, 53-57.
- [5] S.A. Buterin, *On half inverse problem for differential pencils with the spectral parameter in boundary conditions*, Tamkang J. Math. 42 (2011), no. 3, 355-364.
- [6] S.A. Buterin, V.A. Yurko, *Inverse spectral problem for pencils of differential operators on a finite interval*, Vestnik Bashkir. Univ. (2006), no. 4, 8-12 (in Russian).
- [7] S.A. Buterin, V.A. Yurko, *Inverse problems for second-order differential pencils with Dirichlet boundary conditions*. J. Inverse Ill-Posed Probl. 20 (2012), no. 5–6, 855-881.
- [8] D. Chelkak, E. Korotyaev, *Weyl-Titchmarsh functions of vector-valued Sturm-Liouville operators on the unit interval*, J. Func. Anal. 257 (2009), 1546-1588.
- [9] G. Freiling, V. Yurko, *Inverse Sturm-Liouville problems and their applications*, Huntington, NY: Nova Science Publishers, 2001, 305 pp.
- [10] M.G. Gasymov, G.Sh. Gusejnov, *Determination of diffusion operator from spectral data*, Akad. Nauk Azerb. SSR. Dokl. 37 (1981), 19-23.
- [11] R. Hryniv, N. Pronska, *Inverse spectral problems for energy-dependent Sturm-Liouville equations*, Inverse Problems 28 (2012), 085008 (21 pp).
- [12] M.Yu. Ignatiev, V.A. Yurko, *Numerical methods for solving inverse Sturm–Liouville problems*, Results in Mathematics 52 (2008), no. 1–2, 63-74.
- [13] M. Jaulent, C. Jean, *The inverse s-wave scattering problem for a class of potentials depending on energy*, Comm. Math. Phys. 28 (1972), 177-220.
- [14] M. V. Keldysh, *On eigenvalues and eigenfunctions of some classes of nonselfadjoint equations*. Dokl. Akad. Nauk. SSSR 77 (1951), 11-14.
- [15] M.V. Keldysh, *On the completeness of the eigenfunctions of some classes of nonselfadjoint linear operators*, Russian Math. Surveys 26 (1971), no. 4, 15-44.
- [16] A.G. Kostyuchenko, A.A. Shkalikov, *Selfadjoint quadratic operator pencils and elliptic problems*, Funk. Anal. i Prilozhen. 17 (1983), no. 2, 38-61; English transl.: Funct. Anal. Appl. 17 (1983), 109-128.
- [17] V.A. Marchenko, *Sturm-Liouville operators and their applications*. Naukova Dumka, Kiev, 1977 (in Russian); English transl., Birkhauser, 1986.
- [18] A.S. Markus, *Introduction to the spectral theory of polynomial operator pencils*, Translations of Mathematical Monographs 71. American Mathematical Society, Providence, RI, 1988. Translated from the Russian edition (“Shtiintsa”, Kishinev, 1986).

- [19] Ya.V. Mykytyuk, N.S. Trush, *Inverse spectral problems for Sturm-Liouville operators with matrix-valued potentials*, Inverse Problems, 26 (2010), 015009.
- [20] N. Pronska, *Reconstruction of energy-dependent Sturm-Liouville equations from two spectra*, Integral Equations and Operator Theory 76 (2013), no. 3, 403-419.
- [21] A.A. Shkalikov, *Boundary value problems for ordinary differential equations with a parameter in the boundary conditions*, Trudy Sem. Petrovsk. 9 (1983), 190-229 (in Russian).
- [22] M. Yamamoto, *Inverse eigenvalue problem for a vibration of a string with viscous drag*. J. Math. Anal. Appl. 152 (1990), 20-34.
- [23] V.A. Yurko, *On boundary value problems with the parameter in the boundary conditions*, Izvestiya AN SSR. Ser. Matem. 19 (1984), 398-409.
- [24] V.A. Yurko, *An inverse problem for systems of differential equations with nonlinear dependence on the spectral parameter*, Diff. Uravneniya 33 (1997), no. 3, 390-395 (in Russian); English transl. Diff. Equ. 33 (1997), no. 3, 388-394.
- [25] V.A. Yurko, *An inverse problem for pencils of differential operators*, Matem. Sbornik 191 (2000), 137-160.
- [26] V.A. Yurko, *Method of spectral mappings in the inverse problems theory*, Inverse and Ill-posed Problems Series, Utrecht: VSP, 2002.
- [27] V.A. Yurko, *Inverse problems for the matrix Sturm-Liouville equation on a finite interval*, Inverse Problems, 22 (2006), 1139-1149.

Natalia Bondarenko
Department of Mathematics
Saratov State University
Astrakhanskaya 83, Saratov 410026, Russia
E-mail: bondarenkonp@info.sgu.ru

Received: 27.03.2013