



Общероссийский математический портал

Adem Kilicman, Shantha Kumari Kurumujji, Arjun K. Rathie, О новом классе интегралов, включающих обобщенные гипергеометрические функции,
Журн. СФУ. Сер. Матем. и физ., 2024, том 17, выпуск 2, 266–271

<https://www.mathnet.ru/jsfu1156>

Использование Общероссийского математического портала Math-Net.Ru подразумевает, что вы прочитали и согласны с пользовательским соглашением

<https://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 18.97.9.168

21 апреля 2025 г., 20:36:51



EDN: UODEBV
УДК 517.5

On a New Class of Integrals Involving Generalized Hypergeometric Functions

Adem Kilicman*

Department of Mathematics
Institute for Mathematical Research
University Putra Malaysia (UPM)
Selangor, Malaysia

Shantha Kumari Kurumujji†

Department of Mathematics
A J Institute of Engineering and Technology
Visvesvaraya Technological University (VTU), Belagavi
Karnataka, India

Arjun K. Rathie‡

Department of Mathematics
Vedant College of Engineering and Technology
Rajasthan Technical University
Rajasthan State, India

Received 27.04.2023, received in revised form 20.08.2023, accepted 11.01.2024

Abstract. In the theory of hypergeometric and generalized hypergeometric series, classical summation theorems such as those of Gauss, Gauss second, Bailey and Kummer for the series ${}_2F_1$; Watson, Dixon, Whipple and Saalshüz play a key role. Applications of the above mentioned summation theorems are well known. In our present investigation, we aim to evaluate twenty five new class of integrals involving generalized hypergeometric function in the form of a single integral of the form:

$$\int_0^1 x^{c-1}(1-x)^{c-1} {}_3F_2 \left[\begin{matrix} a, b, c + \frac{1}{2} \\ \frac{1}{2}(a+b+i+1), 2c+j \end{matrix} ; 4x(1-x) \right] dx$$

for $i, j = 0, \pm 1, \pm 2$.

The results are established with the help of the generalizations of the classical Watson's summation theorem obtained earlier by Lavoie et al. [2]. Fifty interesting integrals in the form of two integrals (twenty five each) have also been given as special cases of our main findings.

Keywords: generalized hypergeometric function, Watsons theorem, definite integral, beta integral.

Citation: A. Kilicman, S.K. Kumari, A.K. Rathie, On a New Class of Integrals Involving Generalized Hypergeometric Functions, J. Sib. Fed. Univ. Math. Phys., 2024, 17(2), 266–271. EDN: UODEBV.



1. Introduction and preliminaries

The natural generalization of the Gauss's hypergeometric function ${}_2F_1$ is called the generalized hypergeometric function ${}_pF_q$, where $p, q \in \mathbb{N}_0$ defined by [1, 5]

$${}_pF_q \left[\begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} ; z \right] = \sum_{n=0}^{\infty} \frac{(a_1)_n \dots (a_p)_n}{(b_1)_n \dots (b_q)_n} \frac{z^n}{n!} \quad (1)$$

*akilicman@putra.edu.my <https://orcid.org/0000-0002-1217-963X>

†shanthakk99@gmail.co <https://orcid.org/0000-0002-2153-0524>

‡arjunkumarrathie@gmail.com <https://orcid.org/0000-0003-3902-3050>

© Siberian Federal University. All rights reserved

where $(a)_n$ is the well known Pochhammer symbol (or the raised factorial or the shifted factorial since $(1)_n = n!$) defined for any complex $a \in \mathbb{C}$ by

$$\begin{aligned} (a)_n &= \frac{\Gamma(a+n)}{\Gamma(a)}, \quad (a \in \mathbb{C} \setminus \mathbb{Z}_0^-) = \\ &= \begin{cases} a(a+1)\dots(a+n-1), & (n \in \mathbb{N}) \\ 1, & (n = 0) \end{cases} \end{aligned} \quad (2)$$

where Γ is the well known Gamma function.

For a detailed study about hypergeometric and generalized hypergeometric functions, we refer the standard texts [1, 5].

In the theory of hypergeometric and generalized hypergeometric functions, classical summation theorems such as those of Gauss, Gauss second, Kummer and Bailey for the series ${}_2F_1$; Watson, Dixon, Whipple and Saalschütz for the series ${}_3F_2$ play a key role.

Later, the above mentioned classical summation theorems have been generalized by Lavoie et al. [2-4].

However, in our present investigation, we are interested in the following classical Watson's summation theorem [1].

$${}_3F_2 \left[\begin{matrix} a, b, c \\ \frac{1}{2}(a+b+1), 2c \end{matrix}; 1 \right] = \frac{\Gamma(\frac{1}{2}) \Gamma(c + \frac{1}{2}) \Gamma(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}) \Gamma(c - \frac{1}{2}a - \frac{1}{2}b + \frac{1}{2})}{\Gamma(\frac{1}{2}a + \frac{1}{2}) \Gamma(\frac{1}{2}b + \frac{1}{2}) \Gamma(c - \frac{1}{2}a + \frac{1}{2}) \Gamma(c - \frac{1}{2}b + \frac{1}{2})} \quad (3)$$

provided $\Re(2c - a - b) > -1$, and its following generalization due to Lavoie et al. [2].

$$\begin{aligned} &{}_3F_2 \left[\begin{matrix} a, b, c \\ \frac{1}{2}(a+b+i+1), 2c+j \end{matrix}; 1 \right] = \\ &= \mathcal{A}_{i,j} \frac{2^{a+b+i-2} \Gamma(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}i + \frac{1}{2}) \Gamma(c + [\frac{j}{2}] + \frac{1}{2}) \Gamma(c - \frac{1}{2}(a+b+|i+j|-j-1))}{\Gamma(\frac{1}{2}) \Gamma(a) \Gamma(b)} \times \\ &\times \left\{ \mathcal{B}_{i,j} \frac{\Gamma(\frac{1}{2}a + \frac{1}{4}(1 - (-1)^i)) \Gamma(\frac{1}{2}b)}{\Gamma(c - \frac{1}{2}a + \frac{1}{2} + [\frac{j}{2}] - \frac{1}{4}(-1)^j (1 - (-1)^i)) \Gamma(c - \frac{1}{2}b + \frac{1}{2} + [\frac{j}{2}])} + \right. \\ &\left. + \mathcal{C}_{i,j} \frac{\Gamma(\frac{1}{2}a + \frac{1}{4}(1 + (-1)^i)) \Gamma(\frac{1}{2}b + \frac{1}{2})}{\Gamma(c - \frac{1}{2}a + [\frac{j+1}{2}] + \frac{1}{4}(-1)^j (1 - (-1)^i)) \Gamma(c - \frac{1}{2}b + [\frac{j+1}{2}])} \right\} = \\ &= \Omega \quad (\text{let}) \end{aligned} \quad (4)$$

for $i, j = 0, \pm 1, \pm 2$.

For $i = j = 0$, the result (4) reduces to classical Watson's summation theorem (3).

Here, $[x]$ denotes the greatest integer less than or equal to x and the modulus is denoted by $|x|$. For the expressions of the coefficients $\mathcal{A}_{i,j}$, $\mathcal{B}_{i,j}$ and $\mathcal{C}_{i,j}$, one can refer [2].

The aim of this paper is to evaluate twenty five integrals involving generalized hypergeometric function in the form of a single integral of the form

$$\int_0^1 x^{c-1} (1-x)^{c-1} {}_3F_2 \left[\begin{matrix} a, b, c + \frac{1}{2} \\ \frac{1}{2}(a+b+i+1), 2c+j \end{matrix}; 4x(1-x) \right] dx$$

for $i, j = 0, \pm 1, \pm 2$.

The results are derived with the help of generalized Watson's summation theorem on the sum of a ${}_3F_2$ given by (4). Fifty interesting integrals in the form of two integrals (twenty five each) have also been given as special cases of our main findings.

2. Main integrals

The twenty five integrals in the form of a single integral to be evaluated in this paper is given in the following theorem.

Theorem 2.1. For $\Re(c) > 0$, $\Re(2c - a - b + i + 2j + 1) > 0$, for $i, j = 0, \pm 1, \pm 2$, the following integral formula holds.

$$\int_0^1 x^{c-1} (1-x)^{c-1} {}_3F_2 \left[\begin{matrix} a, b, c + \frac{1}{2} \\ \frac{1}{2}(a+b+i+1), 2c+j \end{matrix}; 4x(1-x) \right] dx = \frac{\Gamma(c) \Gamma(c)}{\Gamma(2c)} \Omega \quad (5)$$

where Ω is the same as given in (4).

Proof: The proof of our theorem is quite straight forward. For this, we proceed as follows. Denoting the left hand side of (5) by I , we have

$$I = \int_0^1 x^{c-1} (1-x)^{c-1} {}_3F_2 \left[\begin{matrix} a, b, c + \frac{1}{2} \\ \frac{1}{2}(a+b+i+1), 2c+j \end{matrix}; 4x(1-x) \right] dx \quad (6)$$

Now expressing ${}_3F_2$ as a series, changing the order of integration and summation, which is easily seen to be justified due to the uniform convergence of the series in the interval $(0, 1)$, we have

$$I = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n (c)_n 2^{2n}}{\left(\frac{1}{2}(a+b+i+1)\right)_n (2c+j)_n} \int_0^1 x^{c+n-1} (1-x)^{c+n-1} dx \quad (7)$$

Evaluating the Beta integral and using the result

$$(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)}$$

we have, after some simplification

$$I = \frac{\Gamma(c) \Gamma(c)}{\Gamma(2c)} \sum_{n=0}^{\infty} \frac{(a)_n (b)_n (c)_n}{\left(\frac{1}{2}(a+b+i+1)\right)_n (2c+j)_n n!} \quad (8)$$

Now summing up the series, we have

$$I = \frac{\Gamma(c) \Gamma(c)}{\Gamma(2c)} = {}_3F_2 \left[\begin{matrix} a, b, c \\ \frac{1}{2}(a+b+i+1), 2c+j \end{matrix}; 4x(1-x) \right] \quad (9)$$

We now observe that the ${}_3F_2$ appearing can be evaluated with the help of known result (4) and we easily arrive at the right hand side of (5).

This completes the proof of the theorem. \square

3. Special cases

In this section, we shall mention a large number of very interesting special cases of our main findings.

For this, we observe here that, if in (5), we let $b = -2n$ and replace a by $a + 2n$ or we let $b = -2n - 1$ and replace a by $a + 2n + 1$. In each case, one of the two terms appearing on the right-hand side of (5) will vanish and we get fifty interesting special cases (twenty five each) given below in the form of two corollaries.

Corollary 3.1. For $i, j = 0, \pm 1, \pm 2$, the following twenty five results holds.

$$\int_0^1 x^{c-1}(1-x)^{c-1} {}_3F_2 \left[\begin{matrix} -2n, a+2n, c+\frac{1}{2} \\ \frac{1}{2}(a+i+1), 2c+j \end{matrix}; 4x(1-x) \right] dx =$$

$$= D_{i,j} \frac{\Gamma(c) \Gamma(c)}{\Gamma(2c)} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}a - c + \frac{3}{4} - \frac{(-1)^i}{4} - \left[\frac{1}{2}j + \frac{1}{4}(1 + (-1)^i)\right]\right)_n}{\left(c + \frac{1}{2} + \left[\frac{j}{2}\right]\right)_n \left(\frac{1}{2}a + \frac{1}{4}(1 + (-1)^i)\right)_n} \tag{10}$$

where the coefficients $D_{i,j}$ are as in Tables (1) and (2) given below.

Table 1. Table for $D_{i,j}$, $i = 0, \pm 1, \pm 2$ and $j = -2, -1, 0$

$i \setminus j$	-2	-1	0
2	$\frac{(a+1)[(c-1)(a-1)+2n(a+2n)]}{(c-1)(a+4n-1)(a+4n+1)}$	$\frac{(a+1)(a-1)}{(a+4n+1)(a+4n-1)}$	$\frac{(a+1)[(a-1)(2c-a-1)-4n(a+2n)]}{(2c-a-1)(a+4n+1)(a+4n-1)}$
1	$\frac{a(c+2n-1)}{(c-1)(a+4n)}$	$\frac{a}{a+4n}$	$\frac{a}{a+4n}$
0	$1 - \frac{2n(a+2n)}{(c-1)(2c-a-3)}$	1	1
-1	$1 - \frac{2n(2c+a+4n-2)}{(c-1)(2c-a-4)}$	$1 - \frac{4n}{(2c-a-2)}$	1
-2	$D_{-2,-2}$	$1 - \frac{8n(a+2n)}{(a-1)(2c-a-3)}$	$1 - \frac{4n(a+2n)}{(a-1)(2c-a-1)}$

$$D_{-2,-2} = 1 - \frac{2an(6c+a-7)(2c-a-3) - 4n^2[5a^2 - 4a - 21 - 4c(3c-a-8)] - 64n^3(a+n)}{(c-1)(a-1)(2c-a-3)(2c-a-5)}$$

Table 2. Table for $D_{i,j}$, $i = 0, \pm 1, \pm 2$ and $j = 1, 2$

$i \setminus j$	1	2
2	$\frac{(a+1)[(a-1)(2c-a-1)-8n(a+2n)]}{(2c-a-1)(a+4n+1)(a+4n-1)}$	$D_{2,2}$
1	$\frac{a(2c-a-4n)}{(2c-a)(a+4n)}$	$\frac{a[(c+1)(2c-a)-2n(2c+a+4n+2)]}{(c+1)(2c-a)(a+4n)}$
0	1	$1 - \frac{2n(a+2n)}{(c+1)(2c-a+1)}$
-1	1	$1 + \frac{2n}{(c+1)}$
-2	1	$1 + \frac{2n(a+2n)}{(c+1)(a-1)}$

$$D_{2,2} = \frac{(a+1) \left((a-1)(c+1)(2c-a+1)(2c-a-1) - 2an(6c+a+5)(2c-a+1) + 4n^2(5a^2+4a-5-4c(3c-a+4)) + 64n^3(a+n) \right)}{(c+1)(2c-a+1)(2c-a-1)(a+4n+1)(a+4n-1)}$$

Corollary 3.2. For $i, j = 0, \pm 1, \pm 2$, the following twenty five results holds.

$$\int_0^1 x^{c-1}(1-x)^{c-1} {}_3F_2 \left[\begin{matrix} -2n-1, a+2n+1, c+\frac{1}{2} \\ \frac{1}{2}(a+i+1), 2c+j \end{matrix}; 4x(1-x) \right] dx =$$

$$= E_{i,j} \frac{\Gamma(c) \Gamma(c)}{\Gamma(2c)} \frac{\left(\frac{3}{2}\right)_n \left(\frac{1}{2}a - c + \frac{5}{4} + \frac{(-1)^i}{4} - \left[\frac{1}{2}j + \frac{1}{4}(1 + (-1)^i)\right]\right)_n}{\left(c + \frac{1}{2} + \left[\frac{j+1}{2}\right]\right)_n \left(\frac{1}{2}a + \frac{1}{4}(3 - (-1)^i)\right)_n} \tag{11}$$

where the coefficient $E_{i,j}$ are as in Tables (3) and (4) given below.

Table 3. Table for $E_{i,j}$ $i = 0, \pm 1, \pm 2$ and $j = -2, -1, 0$

$i \setminus j$	-2	-1	0
2	$\frac{(a+1)(2c-a-3)}{(c-1)(a+4n+1)(a+4n+3)}$	$\frac{(a+1)(4c-a-3)}{(a+4n+1)(a+4n+3)(2c-1)}$	$\frac{2(a+1)}{(a+4n+1)(a+4n+3)}$
1	$\frac{(c-a-2n-2)}{(c-1)(a+4n+2)}$	$\frac{2c-a-2}{(a+4n+2)(2c-1)}$	$\frac{1}{a+4n+2}$
0	$\frac{-1}{(c-1)}$	$\frac{-1}{(2c-1)}$	0
-1	$E_{-1,-2}$	$\frac{-(2c+a+4n)}{a(2c-1)}$	$\frac{-1}{a}$
-2	$\frac{-(2c+a+4n-1)(2c-a-4n-5)}{(a-1)(c-1)(2c-a-5)}$	$E_{-2,-1}$	$\frac{-2}{(a-1)}$

$$E_{-2,-1} = - \frac{[(4c+a-1)(2c-a-3) - 8n(a+2n+2)]}{(a-1)(2c-1)(2c-a-3)}$$

$$E_{-1,-2} = - \frac{[(c+a)(2c-a-4) - 2n(3a-2c+4n+6)]}{a(c-1)(2c-a-4)}$$

Table 4. Table for $E_{i,j}$ $i = 0, \pm 1, \pm 2$ and $j = 1, 2$

$i \setminus j$	1	2
2	$E_{2,1}$	$\frac{(a+1)(2c+a+4n+3)(2c-a-4n-1)}{(c+1)(2c-a-1)(a+4n+1)(a+4n+3)}$
1	$\frac{(2c+a+4n+2)}{(2c+1)(a+4n+2)}$	$\frac{(c+a+2)(2c-a)-2n(3a-2c+4n+2)}{(c+1)(2c-a)(a+4n+2)}$
0	$\frac{1}{(2c+1)}$	$\frac{1}{(c+1)}$
-1	$\frac{-(2c-a)}{a(2c+1)}$	$\frac{-(c-a-2n)}{a(c+1)}$
-2	$\frac{-(4c-a+1)}{(a-1)(2c+1)}$	$\frac{-(2c-a+1)}{(a-1)(c+1)}$

$$E_{2,1} = \frac{(a+1)[(4c+a+3)(2c-a-1) - 8n(a+2n+2)]}{(a+4n+1)(a+4n+3)(2c+1)(2c-a-1)}$$

In particular, in (10), if we take $i = j = 0$, we get the following interesting result.

$$\int_0^1 x^{c-1}(1-x)^{c-1} {}_3F_2 \left[\begin{matrix} -2n, a+2n, c+\frac{1}{2} \\ \frac{1}{2}(a+b+1), 2c \end{matrix}; 4x(1-x) \right] dx =$$

$$= \frac{\Gamma(c) \Gamma(c)}{\Gamma(2c)} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}a - c + \frac{1}{2}\right)_n}{\left(c + \frac{1}{2}\right)_n \left(\frac{1}{2}a + \frac{1}{2}\right)_n} \quad (12)$$

Similarly, in (11), if we take $i = j = 0$, we get the following elegant result.

$$\int_0^1 x^{c-1}(1-x)^{c-1} {}_3F_2 \left[\begin{matrix} -2n-1, a+2n+1, c+\frac{1}{2} \\ \frac{1}{2}(a+b+1), 2c \end{matrix}; 4x(1-x) \right] dx = 0 \quad (13)$$

Similarly, we can obtain other results. We, however, prefer to omit the details.

Conclusions

In this paper, we have evaluated twenty five interesting integrals involving generalized hypergeometric function in the form of a single integral.

The results are established with the help of generalization of classical Watson's summation theorem obtained earlier by Lavoie et al. [2].

Fifty interesting integrals in the form of two integrals (twenty five each) have also been evaluated as special cases of our main findings.

We conclude this paper by remarking that the interesting applications of the integrals obtained in this paper are under investigations and will be published soon.

References

- [1] W.N.Bailey, Generalized Hypergeometric Series, Cambridge Tracts in Mathematics and Mathematical Physics, no. 32, Stechert-Hafner, New York, NY, USA, 1964.
- [2] J.L.Lavoie, F.Groncin, A.K.Rathie, Generalizations of Watson's theorem on the sum of a ${}_3F_2$, *Indian J. Math.*, **34**(1992), no. 2, 23–32.
- [3] J.L.Lavoie, F.Groncin, A.K.Rathie, K.Arora, Generalizations of Dixon's theorem on the sum of a ${}_3F_2$, *Math. Comput.*, **62**(1994), 267–276, .
- [4] J.L.Lavoie, F.Groncin, A.K.Rathie, Generalizations of Whipple's theorem on the sum of a ${}_3F_2$, *J. Comput. Appl. Math.*, **72**(1996), 293–300.
- [5] E.D.Rainville, *Special Functions*, Macmillan Company, New York, 1960; Reprinted by Chelsea Publishing Company, Bronx, New York, 1971.

О новом классе интегралов, включающих обобщенные гипергеометрические функции

Адем Киликман

Институт математических исследований
Университет Путра Малайзия (UPM)
Селангор, Малайзия

Шанта Кумари Курумуджи

АИ Институт инженерии и технологий
Висвесварайский технологический университет (VTU), Белагави
Карнатака, Индия

Арджун К. Рати

Ведантский инженерно-технологический колледж
Технический университет Раджастана
Штат Раджастан, Индия

Аннотация. В теории гипергеометрических и обобщенных гипергеометрических функций классические теоремы суммирования, такие как теоремы Гаусса, Бейли и Каммера для серии ${}_2F_1$; Уотсона, Диксона, Уиппла и Саалшуз, играют ключевую роль. Приложения вышеупомянутых теорем о суммировании хорошо известны. В нашем настоящем исследовании мы стремимся оценить двадцать пять новых классов интегралов, включающих обобщенную гипергеометрическую функцию в форме единого интеграла:

$$\int_0^1 x^{c-1}(1-x)^{c-1} {}_3F_2 \left[\begin{matrix} a, b, c + \frac{1}{2} \\ \frac{1}{2}(a+b+i+1), 2c+j \end{matrix}; 4x(1-x) \right] dx$$

for $i, j = 0, \pm 1, \pm 2$.

Результаты устанавливаются с помощью обобщений теоремы классической суммы Уотсона, полученной ранее Лавойе и др. [2]. Пятьдесят интересных интегралов в форме двух видов интегралов (двадцать пять каждый) также были даны в качестве особых случаев наших основных результатов.

Ключевые слова: обобщенная гипергеометрическая функция, теорема Уотсона, определенный интеграл, бета-интеграл.