



# Math-Net.Ru

Общероссийский математический портал

Natalia P. Bondarenko, Асимптотики спектральных данных краевых задач четвертого порядка, *Известия Иркутского государственного университета. Серия Математика*, 2024, том 47, 31–46

DOI: 10.26516/1997-7670.2024.47.31

Использование Общероссийского математического портала Math-Net.Ru подразумевает, что вы прочитали и согласны с пользовательским соглашением  
<http://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 18.97.14.90

7 февраля 2025 г., 17:19:42





Серия «Математика»  
2024. Т. 47. С. 31–46

Онлайн-доступ к журналу:  
<http://mathizv.isu.ru>

---

---

ИЗВЕСТИЯ  
Иркутского  
государственного  
университета

---

---

Research article

УДК 517.984

MSC 34L20, 34B09, 34B05, 34E05, 46F10

DOI <https://doi.org/10.26516/1997-7670.2024.47.31>

## Spectral Data Asymptotics for Fourth-Order Boundary Value Problems

Natalia P. Bondarenko<sup>1</sup>✉

<sup>1</sup> Saratov State University, Saratov, Russian Federation

✉ [bondarenkonp@sgu.ru](mailto:bondarenkonp@sgu.ru)

**Abstract.** In this paper, we derive sharp asymptotics for the spectral data (eigenvalues and weight numbers) of the fourth-order linear differential equation with a distribution coefficient and three types of separated boundary conditions. Our methods rely on the recent results concerning regularization and asymptotic analysis for higher-order differential operators with distribution coefficients. The results of this paper have applications to the theory of inverse spectral problems as well as a separate significance.

**Keywords:** fourth-order differential operators, distribution coefficients, eigenvalue asymptotics, weight numbers

**Acknowledgements:** The study was financially supported by the Russian Science Foundation (Project No. 21-71-10001), <https://rscf.ru/en/project/21-71-10001/>

**For citation:** Bondarenko N. P. Spectral Data Asymptotics for Fourth-Order Boundary Value Problems. *The Bulletin of Irkutsk State University. Series Mathematics*, 2024, vol. 47, pp. 31–46.

<https://doi.org/10.26516/1997-7670.2024.47.31>

Научная статья

Асимптотики спектральных данных краевых задач четвертого порядка

Н. П. Бондаренко<sup>1</sup>

<sup>1</sup> Саратовский государственный университет, Саратов, Российская Федерация

✉ [bondarenkonp@sgu.ru](mailto:bondarenkonp@sgu.ru)

**Аннотация.** В работе получены точные асимптотики спектральных данных (собственных значений и весовых чисел) линейного дифференциального уравнения четвертого порядка с коэффициентом-распределением и тремя типами распадающихся краевых условий. Методы исследования опираются на недавние результаты, касающиеся регуляризации и асимптотического анализа для дифференциальных операторов высших порядков с коэффициентами-распределениями. Результаты исследования имеют приложения в теории обратных спектральных задач, а также самостоятельное значение.

**Ключевые слова:** дифференциальные операторы четвертого порядка, коэффициенты-распределения, асимптотики собственных значений, весовые числа

**Благодарности:** Работа выполнена при финансовой поддержке РНФ (проект 21-71-10001), <https://rscf.ru/project/21-71-10001/>

**Ссылка для цитирования:** Bondarenko N. P. Spectral Data Asymptotics for Fourth-Order Boundary Value Problems // Известия Иркутского государственного университета. Серия Математика. 2024. Т. 47. С. 31–46. <https://doi.org/10.26516/1997-7670.2024.47.31>

## 1. Introduction

Consider the differential equation

$$y^{(4)} + (\tau_2(x)y')' + (\tau_1(x)y)' + \tau_1(x)y' + \tau_0(x)y = \lambda y(x), \quad x \in (0, 1), \quad (1.1)$$

where  $\tau_2 \in W_2^1[0, 1]$  and  $\tau_1 \in L_2[0, 1]$  are complex-valued functions,  $\tau_0$  is the generalized function of class  $W_2^{-1}[0, 1]$ , that is,  $\tau_0 = r'_0$ ,  $r_0$  is a complex-valued function of  $L_2[0, 1]$ ,  $\lambda$  is the spectral parameter. We understand equation (1.1) with a generalized function coefficient in terms of the regularization approach of Mirzoev and Shkalikov [15] (see the details in Section 2).

This paper aims to derive sharp asymptotics for the eigenvalues and the weight numbers of the boundary value problems  $\mathcal{L}_k$ ,  $k = 1, 2, 3$ , for equation (1.1) with the boundary conditions

$$\mathcal{L}_1: \quad y(0) = 0, \quad y(1) = y'(1) = y''(1) = 0, \quad (1.2)$$

$$\mathcal{L}_2: \quad y(0) = y'(0) = 0, \quad y(1) = y'(1) = 0, \quad (1.3)$$

$$\mathcal{L}_3: \quad y(0) = y'(0) = y''(0) = 0, \quad y(1) = 0. \quad (1.4)$$

A general effective method for obtaining eigenvalue asymptotics for arbitrary order differential operators is described in the book of Naimark [17]. However, derivation of sharp asymptotics for specific classes of operators requires additional efforts. In recent years, eigenvalue asymptotics for the fourth-order differential operators with various types of boundary conditions were obtained by Badanin and Korotyaev [2;3], Möller and Zinsou [16],

Aliyev et al [1], Polyakov [19–21] (see also the bibliography in the mentioned papers). Motivation for investigation of spectral properties for the fourth-order operators arises from applications in mechanics, geophysics, and other fields (see, e.g., [4; 10; 18]). Anyway, the mentioned studies deal with the case of regular (integrable) coefficients. There are much less results on spectral data asymptotics for differential equations with distribution coefficients belonging to spaces of generalized functions. Mikhailets and Molyboga [13;14] investigated the eigenvalue asymptotics for the even-order differential operators  $\frac{d^{2m}}{dx^{2m}} + q(x)$  with distribution potential  $q \in W_2^{-m}[0, 1]$  and semi-periodic boundary conditions. In [5], spectral data asymptotics have been obtained for higher-order differential operators with distribution coefficients in the general form with separated boundary conditions.

In this paper, relying on the approach and on the results of [5], we derive sharp asymptotics for the eigenvalues of the boundary value problems  $\mathcal{L}_k$ ,  $k = 1, 2, 3$ . In addition, we study the asymptotic behavior of the weight numbers, which are used in the inverse spectral theory together with eigenvalues for recovering higher-order differential operators (see, e.g., [8; 9; 11; 24]). We get explicit formulas for all the constants in the obtained asymptotics in terms of the functions  $\tau_1$  and  $\tau_2$  from equation (1.1). Note that the boundary conditions of the problems  $\mathcal{L}_1$  and  $\mathcal{L}_3$  are non-self-adjoint. Therefore, it is difficult to find the starting index for numbering of their eigenvalues. The results of this paper play an important role in the spectral data characterization for equation (1.1) (see [9]) and also have a separate significance.

## 2. Regularization

In this section, we discuss the regularization of equation (1.1) basing on the results of [6; 15; 23].

Let  $\sigma_0(x)$  and  $\sigma_1(x)$  be the unique functions of  $W_2^1[0, 1]$  such that

$$\sigma_0'' = \tau_0, \quad \sigma_0(0) = \sigma_0(1) = 0, \quad \sigma_1' = \tau_1, \quad \sigma_1(0) = 0. \quad (2.1)$$

Define the matrix-function

$$F(x) = [f_{k,j}(x)]_{k,j=1}^4 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -(\sigma_1 + \sigma_0) & 0 & 1 & 0 \\ 0 & -\tau_2 + 2\sigma_0 & 0 & 1 \\ \sigma_0^2 - \sigma_1^2 & 0 & \sigma_1 - \sigma_0 & 0 \end{bmatrix}, \quad (2.2)$$

the quasi-derivatives

$$y^{[0]} := y, \quad y^{[k]} := (y^{[k-1]})' - \sum_{j=1}^k f_{k,j} y^{[j-1]}, \quad k = \overline{1, 4}, \quad (2.3)$$

and the domain

$$\mathcal{D}_F := \{y: y^{[k]} \in AC[0, 1], k = \overline{0, 3}\}.$$

Note that, for  $y \in \mathcal{D}_F$ , the quasi-derivative  $y^{[4]}$  is correctly defined and integrable on  $(0, 1)$ .

**Proposition 1.** *For any  $y \in \mathcal{D}_F$ , the generalized function  $\ell(y) := y^{(4)} + (\tau_2 y')' + (\tau_1 y)' + \tau_1 y' + \tau_0 y$  is regular and  $\ell(y) = y^{[4]}$ .*

Proposition 1 is a special case of Theorem 2.2 in [6], which was obtained from the results of Vladimirov [23]. It follows from Proposition 1 that, for  $y \in \mathcal{D}_F$ , equation (1.1) can be equivalently represented as the first-order system

$$\vec{y}'(x) = (F(x) + \Lambda)\vec{y}(x), \quad x \in (0, 1), \quad (2.4)$$

where

$$\vec{y}(x) = \begin{bmatrix} y(x) \\ y^{[1]}(x) \\ y^{[2]}(x) \\ y^{[3]}(x) \end{bmatrix}, \quad \Lambda = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \lambda & 0 & 0 & 0 \end{bmatrix}.$$

Indeed, the first three rows of the system (2.4) coincide with the definitions of the quasi-derivatives (2.3), and the last row is equivalent to the equation  $y^{[4]} = \lambda y$ . Thus, we have regularized equation (1.1) by the reduction to the system (2.4) with the integrable matrix function  $F(x)$ .

Note that, for regularization of equation (1.1), different constructions of an associated matrix  $F(x)$  can be used (see [6; 15]). Anyway, it has been proved in [7] that the choice of an associated matrix does not influence the spectral data, which will be considered in the next sections. For the purposes of this paper, it is convenient to use the associated matrix (2.2), because it contains all the coefficients  $\tau_2$ ,  $\sigma_1$ , and  $\sigma_0$  of the same smoothness on the same diagonal.

### 3. Birkhoff-type solutions

In this section, we construct the Birkhoff-type solutions of the system (2.4) by the well-known method (see, e.g., [17; 22]), using the technique of [25]. Consequently, we get the Birkhoff-type solutions for equation (1.1).

In the asymptotic estimates below, we use the following **notations**:

- Put  $\lambda = \rho^4$  and divide the complex  $\rho$ -plane into the sectors

$$\Gamma_\kappa := \left\{ \rho \in \mathbb{C}: \frac{\pi(\kappa - 1)}{8} < \arg \rho < \frac{\pi\kappa}{8} \right\}, \quad \kappa = \overline{1, 8}.$$

- For  $\kappa \in \{1, 2, \dots, 8\}$ ,  $h > 0$ , and  $\rho^* > 0$ , introduce the extended sector

$$\Gamma_{\kappa, h, \rho^*} := \left\{ \rho \in \mathbb{C} : \rho + h \exp\left(\frac{i\pi(\kappa-1/2)}{8}\right) \in \Gamma_{\kappa}, |\rho| > \rho^* \right\}.$$

- For a fixed sector  $\Gamma_{\kappa}$ , denote by  $\{\omega_k\}_{k=1}^4$  the roots of the equation  $\omega^4 = 1$  numbered so that

$$\operatorname{Re}(\rho\omega_1) < \operatorname{Re}(\rho\omega_2) < \operatorname{Re}(\rho\omega_3) < \operatorname{Re}(\rho\omega_4), \quad \rho \in \Gamma_{\kappa}. \quad (3.1)$$

Put  $\Omega := [\omega_k^{j-1}]_{j,k=1}^4$ .

- The notation  $\Upsilon(\rho)$  is used for various scalar and vector functions such that

$$\Upsilon(\rho) \rightarrow 0 \quad \text{as} \quad |\rho| \rightarrow \infty, \quad \rho \in \bar{\Gamma}_{\kappa, h, \rho^*}, \quad (3.2)$$

$$\{\|\Upsilon(\rho_n)\|\} \in l_2 \quad \text{for any non-condensing sequence } \{\rho_n\} \in \Gamma_{\kappa, h, \rho^*}, \quad (3.3)$$

where  $\|\cdot\|$  stands for the vector norm if  $\Upsilon(\rho)$  is a vector. Recall that a sequence  $\{\rho_n\}_{n=1}^{\infty}$  is called *non-condensing* if

$$\sup_{t>0} (N(t+1) - N(t)) < \infty, \quad N(t) := \#\{n \in \mathbb{N} : |\rho_n| \leq t\}.$$

- $\Upsilon(x, \rho)$  stands for various scalar and vector functions having the properties (3.2) and (3.3) for each fixed  $x \in [0, 1]$ .
- $\varepsilon(\rho)$  is used for various functions of form  $\frac{\Upsilon(\rho)}{\rho^2}$ .
- $\{\varkappa_n\}$  stands for various  $l_2$ -sequences.

The change of variables

$$Y(x) = \Omega^{-1} \operatorname{diag} \{1, \rho^{-1}, \rho^{-2}, \rho^{-3}\} \bar{y}(x)$$

transforms the system (2.4) into

$$LY := \frac{1}{\rho} Y' - A(x, \rho) Y = 0, \quad (3.4)$$

where

$$A(x, \rho) = A_{(0)} + \frac{A_{(2)}(x)}{\rho^2} + \frac{A_{(4)}(x)}{\rho^4},$$

$$A_{(0)} = \text{diag}\{\omega_k\}_{k=1}^4, \quad A_{(1)} = A_{(3)} = 0, \quad (3.5)$$

$$A_{(2)} := \Omega^{-1} \begin{bmatrix} 0 & 0 & 0 & 0 \\ -(\sigma_1 + \sigma_0) & 0 & 0 & 0 \\ 0 & -\tau_2 + 2\sigma_0 & 0 & 0 \\ 0 & 0 & \sigma_1 - \sigma_0 & 0 \end{bmatrix} \Omega, \quad (3.6)$$

$$A_{(4)} := \Omega^{-1} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \sigma_0^2 - \sigma_1^2 & 0 & 0 & 0 \end{bmatrix} \Omega.$$

Clearly, the entries of  $A_{(2)}(x)$  and  $A_{(4)}(x)$  belong to  $W_2^1[0, 1]$ .

Below, we suppose that  $\rho \in \bar{\Gamma}_{\kappa, h, \rho^*}$  for some  $\kappa, h > 0, \rho^* > 0$ , and  $\{\omega_k\}_{k=1}^4$  are numbered in the order (3.1) for  $\rho \in \Gamma_{\kappa}$ . Note that, for  $\rho \in \bar{\Gamma}_{\kappa, h, \rho^*}$ , we have  $\text{Re}(\rho\omega_k) - \text{Re}(\rho\omega_{k+1}) \leq c_h < \infty$ .

Let us search for approximate solutions of (3.4) in the form

$$U_k(x, \rho) = \exp(\rho\omega_k x) \left( g_{(0)k} + \frac{g_{(1)k}(x)}{\rho} + \frac{g_{(2)k}(x)}{\rho^2} \right), \quad k = \overline{1, 4}, \quad (3.7)$$

where  $g_{(\mu)k} = [g_{(\mu)k,j}]_{j=1}^4$  are column vectors. Furthermore,  $U_k(x, \rho)$  has to satisfy the estimate

$$LU_k = \exp(\rho\omega_k x) \rho^{-3} (h_k(x) + O(\rho^{-1})), \quad \rho \in \bar{\Gamma}_{\kappa, h, \rho^*}, \quad |\rho| \rightarrow \infty, \quad (3.8)$$

where  $h_k(x) = [h_{k,j}(x)]_{j=1}^4$  is a column vector such that  $h_{k,k} = 0$ , and the  $O$ -estimate is uniform with respect to  $x \in [0, 1]$ .

Due to [25], the vectors  $g_{(\mu)k}$ ,  $\mu = 0, 1, 2$ , and  $h_k$  have to fulfill the relations

$$A_{(0)}g_{(0)k} = \omega_k g_{(0)k},$$

$$A_{(0)}g_{(\mu)k} = \omega_k g_{(\mu)k} + g'_{(\mu-1)k} - \sum_{j=1}^{\mu} A_{(j)}g_{(\mu-j)k}, \quad \mu = 1, 2,$$

$$h_k = g'_{(2)k} - \sum_{j=1}^3 A_{(j)}g_{(3-j)k}, \quad h_{k,k} = 0.$$

Therefore, using (3.5) and (3.6), we obtain

$$g_{(0)k} = e_k, \quad g_{(1)k}(x) = -\frac{1}{4\omega_k} \tau_2^{\langle -1 \rangle}(x) e_k, \quad (3.9)$$

$$g_{(2)k,j}(x) = \begin{cases} \frac{1}{(4\omega_k)^2} \tau_2^{\langle -2 \rangle}(x), & j = k, \\ -\frac{a_{jk}(x)}{\omega_j - \omega_k}, & j \neq k. \end{cases}, \quad (3.10)$$

$$h_{k,j}(x) = -\frac{a'_{jk}(x)}{\omega_j - \omega_k} - \frac{1}{4\omega_k} a_{jk}(x) \tau_2^{\langle -1 \rangle}(x), \quad j \neq k,$$

where  $e_k$  is the  $k$ -th column of the unit matrix,

$$\tau_2^{\langle -1 \rangle}(x) = \int_0^x \tau_2(t) dt, \quad \tau_2^{\langle -2 \rangle}(x) = \int_0^x \tau_2(t) \tau_2^{\langle -1 \rangle}(t) dt$$

and  $[a_{jk}(x)]_{j,k=1}^4 := A_{(2)}(x)$ . Note that  $h_{k,j} \in L_2[0, 1]$ .

Thus, the functions  $U_k(x, \rho)$ ,  $k = \overline{1, 4}$ , given by (3.7) with the coefficients (3.9), (3.10) satisfy (3.8). Let  $U(x, \rho)$  be the matrix function consisting of the columns  $U_k(x, \rho)$  and let  $V_j(x, \rho)$ ,  $j = \overline{1, 4}$ , be the rows of  $V(x, \rho) := (U(x, \rho))^{-1}$ . Denote

$$L^* Z := \frac{1}{\rho} Z' - Z A(x, \rho).$$

For  $k = \overline{1, 4}$ , consider the integral equation

$$\begin{aligned} Y_k(x, \rho) = & U_k(x, \rho) + \rho \int_0^x \left( \sum_{j=1}^k U_j(x, \rho) L^* V_j(t, \rho) \right) Y_k(t, \rho) dt \\ & - \rho \int_x^1 \left( \sum_{j=k+1}^4 U_j(x, \rho) L^* V_j(t, \rho) \right) Y_k(t, \rho) dt, \end{aligned} \quad (3.11)$$

whose solution  $Y_k(x, \rho)$  solves the system (3.4). Changing the variables

$$Y_k(x, \rho) = \exp(\rho \omega_k x) W_k(x, \rho), \quad U_k(x, \rho) = \exp(\rho \omega_k x) W_k^0(x, \rho) \quad (3.12)$$

and using (3.8), we transform equation (3.11) into

$$W_k(x, \rho) = W_k^0(x, \rho) + \frac{1}{\rho^2} \int_0^1 B_k(x, t, \rho) W_k(t, \rho) dt, \quad (3.13)$$

where the function  $B_k(x, t, \rho)$  equals

$$\begin{cases} -\sum_{j=1}^k \exp(\rho(\omega_j - \omega_k)(x-t)) W_j^0(x, \rho) (h_{k,j}(t) + O(\rho^{-1})), & x \geq t, \\ \sum_{j=k+1}^4 \exp(\rho(\omega_j - \omega_k)(x-t)) W_j^0(x, \rho) (h_{k,j}(t) + O(\rho^{-1})), & x < t, \end{cases}$$



and the  $O$ -estimates are uniform with respect to  $x \in [0, 1]$  and  $\rho \in \overline{\Gamma}_{\kappa, h, \rho^*}$  as  $|\rho| \rightarrow \infty$ . The analysis of the integral term in (3.13) shows that

$$W_k(x, \rho) - W_k^0(x, \rho) = \frac{\Upsilon(x, \rho)}{\rho^2}, \quad (3.14)$$

where  $\Upsilon(x, \rho)$  denotes various vector functions satisfying (3.2) and (3.3) for each fixed  $x \in [0, 1]$ .

Combining (3.7), (3.9), (3.10), (3.12), and (3.14) all together, we obtain the fundamental solutions  $\{Y_k(x, \rho)\}_{k=1}^4$  of the system (3.4) for  $\rho \in \overline{\Gamma}_{\kappa, h, \rho^*}$ :

$$Y_k(x, \rho) = \exp(\rho\omega_k x) \left( \left( 1 - \frac{\tau_2^{(-1)}(x)}{4\omega_k \rho} + \frac{\tau_2^{(-2)}(x)}{(4\omega_k \rho)^2} \right) e_k - \sum_{\substack{j=1 \\ j \neq k}}^4 \frac{a_{jk}(x)}{\omega_j - \omega_k} e_j + \frac{\Upsilon(x, \rho)}{\rho^2} \right).$$

Returning from the system (3.4) to equation (1.1), we arrive at the following theorem.

**Theorem 1.** *For any  $\kappa = \overline{1, 8}$ ,  $h > 0$  and some  $\rho^* > 0$ , equation 1.1 has a fundamental system of solutions  $\{y_k(x, \rho)\}_{k=1}^4$  whose quasi-derivatives  $y_k^{[j]}(x, \rho)$ ,  $k = \overline{1, 4}$ ,  $j = \overline{0, 3}$ , are continuous by  $x \in [0, 1]$  and  $\rho \in \overline{\Gamma}_{\kappa, h, \rho^*}$ , analytic in  $\rho \in \Gamma_{\kappa, h, \rho^*}$ , for each fixed  $x \in [0, 1]$ , and satisfy the asymptotic relation*

$$y_k^{[j]}(x, \rho) = (\rho\omega_k)^j \exp(\rho\omega_k x) \left( 1 - \frac{\tau_2^{(-1)}(x)}{4\rho\omega_k} + \frac{\tau_2^{(-2)}(x)}{(4\rho\omega_k)^2} - \sum_{\substack{l=1 \\ l \neq k}}^4 \left( \frac{\omega_l}{\omega_k} \right)^j \frac{a_{lk}(x)}{\rho^2(\omega_l - \omega_k)} + \frac{\Upsilon(x, \rho)}{\rho^2} \right). \quad (3.15)$$

It follows from (2.1) and (3.6) that

$$\begin{aligned} a_{lk}(0) &= -\frac{1}{4}\tau_2(0)\omega_l^{-2}\omega_k, \\ a_{lk}(1) &= -\frac{1}{4}\tau_2(1)\omega_l^{-2}\omega_k + \frac{1}{4}\sigma_1(1)(\omega_l^{-3}\omega_k^2 - \omega_l^{-1}). \end{aligned}$$

Using the latter relations together with (3.15), we conclude that, for  $k = \overline{1, 4}$ ,  $j = \overline{0, 3}$ ,

$$y_k^{[j]}(0, \rho) = (\rho\omega_k)^j \left( 1 + \frac{c_{(0)jk}t_0}{\rho^2} + \varepsilon(\rho) \right), \quad (3.16)$$

$$y_k^{[j]}(1, \rho) = (\rho\omega_k)^j \left( 1 - \frac{\theta}{4\rho\omega_k} + \frac{\theta^2}{32(\rho\omega_k)^2} + \frac{c_{(0)jk}t_1}{\rho^2} + \frac{c_{(1)jk}\sigma}{\rho^2} + \varepsilon(\rho) \right), \quad (3.17)$$

where  $\varepsilon(\rho) = \frac{\Upsilon(\rho)}{\rho^2}$  and, for brevity,

$$\begin{aligned} \theta &:= \int_0^1 \tau_2(x) dx, \quad t_0 := \tau_2(0), \quad t_1 := \tau_2(1), \quad \sigma := \sigma_1(1) = \int_0^1 \tau_1(x) dx, \\ c_{(0)jk} &:= \frac{1}{4} \sum_{\substack{l=1 \\ l \neq k}}^4 \left( \frac{\omega_l}{\omega_k} \right)^j \frac{\omega_l^{-2} \omega_k}{\omega_l - \omega_k}, \quad c_{(1)jk} := -\frac{1}{4} \sum_{\substack{l=1 \\ l \neq k}}^4 \left( \frac{\omega_l}{\omega_k} \right) \frac{\omega_l^{-3} \omega_k^2 - \omega_l^{-1}}{\omega_l - \omega_k}. \end{aligned}$$

#### 4. Eigenvalue asymptotics

In this section, we obtain the eigenvalue asymptotics for the boundary value problems  $\mathcal{L}_k$ ,  $k = 1, 2, 3$ . The results of [5] imply the following proposition.

**Proposition 2.** *Suppose that  $\tau_2, \tau_1 \in L_2[0, 1]$  and  $\tau_0 \in W_2^{-1}[0, 1]$ . Then, for each  $k = 1, 2, 3$ , the spectrum of the boundary value problem  $\mathcal{L}_k$  is a countable set of eigenvalues  $\{\lambda_{n,k}\}_{n \geq 1}$  (counted with multiplicities) such that*

$$\lambda_{n,k} = (-1)^k \left( \frac{\pi}{\sin \frac{\pi k}{4}} \left( n + \chi_{k,0} + \frac{\chi_{k,1}}{n} + \frac{\varkappa_n}{n} \right) \right)^n, \quad n \in \mathbb{N}, \quad (4.1)$$

where  $\chi_{k,0}$  and  $\chi_{k,1}$  are some constants,  $\chi_{k,0}$  does not depend on  $(\tau_0, \tau_1, \tau_2)$ , and  $\chi_{k,1}$  depends only on  $\theta = \int_0^1 \tau_2(x) dx$ . The notation  $\{\varkappa_n\}$  is used for various sequences of  $l_2$ , in particular, in (4.1) and analogous formulas below,  $\varkappa_n$  depends on  $k$ .

For  $k = 2$ , the constants in (4.1) are known from the previous studies [12; 20]:  $\chi_{2,0} = \frac{1}{2}$ ,  $\chi_{2,1} = -\frac{\theta}{4\pi^2}$ . In this section, we find  $\chi_{k,0}$  and  $\chi_{k,1}$  for  $k = 1, 3$  and improve the remainder term of (4.1), taking  $\tau_2 \in W_2^1[0, 1]$  into account. The main result is presented in the following theorem.

**Theorem 2.** *The following asymptotic relations hold for  $n \geq 1$ :*

$$\begin{aligned} \lambda_{n,2\pm 1} &= - \left( \left( \sqrt{2}\pi n + \frac{\pi}{2\sqrt{2}} \right)^4 - \theta \left( \sqrt{2}\pi n + \frac{\pi}{2\sqrt{2}} \right)^2 \right. \\ &\quad \left. - \frac{t_0 + t_1 \mp 4\sigma}{\sqrt{2}} \left( \sqrt{2}\pi n + \frac{\pi}{2\sqrt{2}} \right) + n\varkappa_n \right), \end{aligned} \quad (4.2)$$

$$\lambda_{n,2} = \left( \pi n + \frac{\pi}{2} \right)^4 - \theta \left( \pi n + \frac{\pi}{2} \right)^2 + (t_0 + t_1) \left( \pi n + \frac{\pi}{2} \right) + n\varkappa_n. \quad (4.3)$$

*Proof.* Note that  $\mathcal{L}_1$  is adjoint to the boundary value problem for equation (1.1) with  $(\tau_0, \tau_1, \tau_2)$  replaced by  $(\overline{\tau_0}, -\overline{\tau_1}, \overline{\tau_2})$  subject to the boundary conditions (1.4). Thus, the eigenvalue asymptotics (4.2) for  $\lambda_{n,1}$  and for  $\lambda_{n,3}$

can be easily obtained from one another. Therefore, it is sufficient to prove (4.2) for  $\lambda_{n,3}$ .

Let  $C_k(x, \lambda)$ ,  $k = \overline{1, 4}$ , be the solutions of equation (1.1) under the initial conditions

$$C_k^{[j-1]}(0, \lambda) = \delta_{k,j}, \quad j = \overline{1, 4}, \quad (4.4)$$

where  $\delta_{k,j}$  is the Kronecker delta.

Note that the boundary conditions (1.4) are equivalent to

$$y(0) = y'(0) = y^{[2]}(0) = 0, \quad y(1) = 1.$$

Hence, the eigenvalues of  $\mathcal{L}_3$  coincide with the zeros of the characteristic function

$$\Delta(\lambda) = \begin{vmatrix} C_1(0, \lambda) & C_2(0, \lambda) & C_3(0, \lambda) & C_4(0, \lambda) \\ C_1'(0, \lambda) & C_2'(0, \lambda) & C_3'(0, \lambda) & C_4'(0, \lambda) \\ C_1^{[2]}(0, \lambda) & C_2^{[2]}(0, \lambda) & C_3^{[2]}(0, \lambda) & C_4^{[2]}(0, \lambda) \\ C_1(1, \lambda) & C_2(1, \lambda) & C_3(1, \lambda) & C_4(1, \lambda) \end{vmatrix}.$$

Let  $\lambda = \rho^4$ ,  $\rho \in \Gamma_{\kappa, h, \rho^*}$ , and let  $\{y_k(x, \rho)\}_{k=1}^4$  be the Birkhoff-type solutions from Theorem 1. Then, the both matrix functions  $C(x, \lambda) := [C_k^{[j-1]}(x, \lambda)]_{j,k=1}^4$  and  $Y(x, \rho) := [y_k^{[j-1]}(x, \rho)]_{j,k=1}^4$  are fundamental solutions of the system (2.4). Hence

$$C(x, \lambda) = Y(x, \rho)\mathcal{A}(\rho),$$

where  $\mathcal{A}(\rho) = (Y(0, \rho))^{-1}$ . Consequently,

$$\Delta(\lambda) = D(\rho) \det(\mathcal{A}(\rho)), \quad (4.5)$$

where

$$D(\rho) = \begin{vmatrix} y_1(0, \rho) & y_2(0, \rho) & y_3(0, \rho) & y_4(0, \rho) \\ y_1'(0, \rho) & y_2'(0, \rho) & y_3'(0, \rho) & y_4'(0, \rho) \\ y_1^{[2]}(0, \rho) & y_2^{[2]}(0, \rho) & y_3^{[2]}(0, \rho) & y_4^{[2]}(0, \rho) \\ y_1(1, \rho) & y_2(1, \rho) & y_3(1, \rho) & y_4(1, \rho) \end{vmatrix}. \quad (4.6)$$

It follows from (3.15) that  $\det(\mathcal{A}(\rho)) \neq 0$  for sufficiently large  $|\rho|$ ,  $\rho \in \overline{\Gamma}_{\kappa, h, \rho^*}$ . Therefore, for such values of  $\rho$ , a complex  $\lambda = \rho^4$  is an eigenvalue of the problem  $\mathcal{L}_3$  if and only if  $D(\rho) = 0$ .

One can easily see that the images of the two closed sectors  $\overline{\Gamma}_1$  and  $\overline{\Gamma}_2$  of the  $\rho$ -plane cover the whole  $\lambda$ -plane. Furthermore, the zeros of  $D(\rho)$  for sufficiently large  $|\rho|$  lie in the neighbourhood of their joint boundary  $\arg \rho = \frac{\pi}{4}$ . For definiteness, consider  $\Gamma_1$ , for which  $\omega_1 = -1$ ,  $\omega_2 = i$ ,  $\omega_3 = -i$ ,  $\omega_4 = 1$ . Substituting (3.16) and (3.17) into (4.6) for  $\rho \in \Gamma_{1, h, \rho^*}$ ,

we derive

$$\begin{aligned}
D(\rho) &= \rho^3 \exp(\rho\omega_4)d(\rho), \\
d(\rho) &= r_1(\rho) - r_2(\rho) \exp(\rho(\omega_3 - \omega_4)) + \varepsilon(\rho), \\
r_1(\rho) &= -4i + \frac{i\theta}{\rho} + \frac{2i\sigma}{\rho^2} - \frac{i(t_0 + t_1)}{2\rho^2} - \frac{i\theta^2}{8\rho^2}, \\
r_2(\rho) &= 4 - \frac{i\theta}{\rho} + \frac{2\sigma}{\rho^2} - \frac{t_0 + t_1}{2\rho^2} - \frac{\theta^2}{8\rho^2}.
\end{aligned} \tag{4.7}$$

Consequently, the equation  $D(\rho) = 0$  in  $\Gamma_{1,h,\rho^*}$  can be reduced to the form

$$\exp(\rho(\omega_4 - \omega_3)) = \frac{r_2(\rho)}{r_1(\rho)} + \varepsilon(\rho) = i + \frac{(1+i)\theta}{4\rho} + \frac{i\sigma}{\rho^2} - \frac{i(t_0 + t_1)}{4\rho^2} + \frac{\theta^2}{16\rho^2} + \varepsilon(\rho).$$

Taking the logarithm and using the standard approach, based on Rouché's theorem (see [5]), we find the roots

$$(1+i)\rho_n = 2\pi in + \frac{\pi i}{2} + \frac{(1+i)\theta}{4i\rho_n} + \frac{\sigma}{\rho_n^2} - \frac{t_0 + t_1}{4\rho_n^2} + \varepsilon(\rho_n)$$

for  $n \geq n_0$ , where  $n_0$  is sufficiently large. Recall that  $\varepsilon(\rho) = \frac{\Upsilon(\rho)}{\rho^2}$ , where  $\Upsilon(\rho)$  satisfies (3.2) and (3.3). Therefore,  $\varepsilon(\rho_n) = \frac{\varkappa_n}{n^2}$ ,  $\{\varkappa_n\} \in l_2$ . Finally, we arrive at the relation

$$\begin{aligned}
\rho_n = \exp\left(\frac{\pi i}{4}\right) &\left( \sqrt{2}\pi n + \frac{\pi}{2\sqrt{2}} - \frac{\theta}{4(\sqrt{2}\pi n + \frac{\pi}{2\sqrt{2}})} - \frac{\sigma}{2\sqrt{2}(\pi n)^2} \right. \\
&\left. + \frac{t_0 + t_1}{8\sqrt{2}(\pi n)^2} + \frac{\varkappa_n}{n^2} \right), \quad n \geq n_0. \tag{4.8}
\end{aligned}$$

Computing  $\lambda_n = \rho_n^4$ , we arrive at the asymptotics (4.2) for sufficiently large values of  $n$ . It remains to show that the numbering in (4.2) starts from  $n = 1$ . By virtue of Theorem 1.1 in [5], it is sufficient to prove this for the eigenvalues  $\{\lambda_{n,3}^0\}$  of the boundary value problem  $\mathcal{L}_3^0$  of the same form as  $\mathcal{L}_3$  but with the zero coefficients  $\tau_j^0 = 0$ ,  $j = 0, 1, 2$ . Using (4.5), (4.6), and the Birkhoff solutions  $y_k^0(x, \rho) = \exp(\rho\omega_k x)$ ,  $k = \overline{1, 4}$ , of the equation  $y^{(4)} = \rho^4 y$ , we obtain the relation

$$\Delta^0(\lambda) = \frac{\sinh(\rho) - \sin(\rho)}{2\rho^3}$$

for the characteristic function of  $\mathcal{L}_3^0$ .

Using the Taylor series

$$\Delta^0(\rho^4) = \frac{1}{3!} + \frac{\rho^4}{7!} + \frac{\rho^8}{11!} + \dots,$$

we show that  $\Delta^0(\rho^4) \neq 0$  for  $|\rho| \leq 5$ . It is sufficient to consider  $0 \leq \arg \rho \leq \frac{\pi}{2}$ . For  $0 \leq \arg \rho \leq \frac{\pi}{8}$ ,  $|\rho| \geq 5$ , we have

$$\begin{aligned} 2|\sinh(\rho)| &\geq |\exp(\rho)| - |\exp(-\rho)| \geq \exp(5 \cos \frac{\pi}{8}) - \exp(-5) > 100, \\ 2|\sin(\rho)| &\leq |\exp(i\rho)| + |\exp(-i\rho)| \leq \exp(5 \sin \frac{\pi}{8}) + 1 < 10, \end{aligned}$$

so  $|\sinh(\rho)| > |\sin(\rho)|$  and  $\Delta^0(\rho^4)$  has no zeros in this sector. Analogously  $\Delta^0(\rho^4) \neq 0$  for  $\frac{3\pi}{8} \leq \arg \rho \leq \frac{\pi}{2}$ . Hence, all the zeros of  $\Delta^0(\rho^4)$  lie in the region

$$\mathcal{G} := \{\rho: |\rho| > 5, \frac{\pi}{8} < \arg \rho < \frac{3\pi}{8}\}.$$

Consider the functions

$$\begin{aligned} f(\rho) &:= 2\sinh(\rho) - 2\sin(\rho) = g(\rho) + s(\rho), \\ g(\rho) &:= \exp(\rho) - i\exp(-i\rho), \quad s(\rho) := i\exp(i\rho) - \exp(\rho). \end{aligned}$$

Obviously, in  $\mathcal{G}$ , the function  $g(\rho)$  has the zeros

$$\rho_n^\diamond = \exp\left(\frac{\pi i}{4}\right) \left(\sqrt{2}\pi n + \frac{\pi}{2\sqrt{2}}\right), \quad n \geq 1,$$

which coincide with the main term of (4.8), and

$$|s(\rho)| \leq 2\exp(-5 \sin \frac{\pi}{8}) < 0.3, \quad \rho \in \mathcal{G}.$$

Define the region

$$\mathcal{G}_\delta := \{\rho \in \mathcal{G}: |\rho - \rho_k^\diamond| \geq \delta, k \geq 1\}, \quad \delta := 0.1.$$

For  $\rho = \rho_n^\diamond + z \in \mathcal{G}_\delta$ ,  $|z| \geq \delta$ , we obtain the estimate

$$|g(\rho)| \geq |\exp(\rho_n^\diamond + z)| |1 - \exp(-(1+i)z)| \geq \exp(5 \cos \frac{3\pi}{8}) \frac{|z|}{\sqrt{2}} > 0.4.$$

Therefore, we get  $\frac{|s(\rho)|}{|g(\rho)|} < 1$  in  $\mathcal{G}_\delta$ . Applying Rouché's theorem to  $f(\rho)$  and  $g(\rho)$ , we conclude that  $f(\rho)$  has only the simple zeros  $\{\rho_n^0\}_{n \geq 1}$  in  $\mathcal{G}$  such that  $|\rho_n^0 - \rho_n^\diamond| < 0.1$ , and, obviously, so does  $\Delta^0(\rho^4)$ . This concludes the proof of (4.2).

The asymptotics (4.3) is proved analogously.  $\square$

**Remark 1.** The main part of the asymptotics (4.3) coincides with the one obtained by Polyakov [20] for the case  $\tau_j \in W_1^{j+1}[0, 1]$ ,  $\tau_j(0) = \tau_j(1)$  for  $j = 0, 2$ ,  $\tau_1 \equiv 0$ . Sharp eigenvalue asymptotics for the fourth-order boundary value problems with the boundary conditions (1.2) and (1.4), to the best of the author's knowledge, have not been investigated before, so the asymptotic formula (4.2) is novel even for the case of regular coefficients.

### 5. Weight numbers

In this section, we recall the definition of the weight numbers from [8] and derive for them precise asymptotics, which play an important role in the spectral data characterization (see [9]).

For  $k = 1, 2, 3$ , define the functions

$$\begin{aligned}\Delta_k(\lambda) &:= \det([C_j^{[4-s]}(1, \lambda)]_{s,j=k+1}^4), \\ \Delta_k^+(\lambda) &:= \det([C_j^{[4-s]}(1, \lambda)]_{s=\overline{k+1,4}, j=\overline{k, k+2, 4}}),\end{aligned}$$

where  $C_j(x, \lambda)$ ,  $j = \overline{1, 4}$ , are the solutions of equation (1.1) satisfying the initial conditions (4.4). One can show in the standard way that, for each  $k \in \{1, 2, 3\}$ , the zeros of  $\Delta_k(\lambda)$  coincide with the eigenvalues  $\{\lambda_{n,k}\}_{n \geq 1}$  of the corresponding boundary value problem  $\mathcal{L}_k$ . The weight numbers are defined as follows:

$$\beta_{n,k} := - \operatorname{Res}_{\lambda=\lambda_{n,k}} \frac{\Delta_k^+(\lambda)}{\Delta_k(\lambda)}, \quad n \geq 1, \quad k = 1, 2, 3.$$

According to the asymptotics of Theorem 2, the eigenvalues  $\lambda_{n,k}$  are simple for  $n \geq n_0$  with a sufficiently large  $n_0$ . Hence, we have

$$\beta_{n,k} = - \frac{\Delta_k^+(\lambda_{n,k})}{\frac{d}{d\lambda} \Delta_k(\lambda_{n,k})}, \quad n \geq n_0, \quad k = 1, 2, 3. \quad (5.1)$$

The main result of this section is provided in the following theorem.

**Theorem 3.** *For  $n \geq n_0$ , the weight numbers satisfy the asymptotic relations*

$$\beta_{n,2\pm 1} = -4\lambda_{n,2\pm 1} \left( 1 + \frac{t_0 + \theta}{8(\pi n)^2} + \frac{\varkappa_n}{n^2} \right), \quad (5.2)$$

$$\beta_{n,2} = -4\lambda_{n,2} \left( 1 + \frac{t_0 + 2\theta}{4(\pi n)^2} + \frac{\varkappa_n}{n^2} \right). \quad (5.3)$$

*Proof.* Similarly to Theorem 2, let us focus on the proof of 5.2 for  $\beta_{n,3}$ . The proofs for  $\beta_{n,1}$  and  $\beta_{n,2}$  are analogous.

Consider  $\rho \in \Gamma_{1,h,\rho^*}$ ,  $\lambda = \rho^4$ . For brevity, put  $\Delta(\lambda) := \Delta_3(\lambda)$  and

$$\Delta^+(\lambda) := \Delta_3^+(\lambda) = - \begin{vmatrix} C_1(0, \lambda) & C_2(0, \lambda) & C_3(0, \lambda) & C_4(0, \lambda) \\ C_1'(0, \lambda) & C_2'(0, \lambda) & C_3'(0, \lambda) & C_4'(0, \lambda) \\ C_1^{[3]}(0, \lambda) & C_2^{[3]}(0, \lambda) & C_3^{[3]}(0, \lambda) & C_4^{[3]}(0, \lambda) \\ C_1(1, \lambda) & C_2(1, \lambda) & C_3(1, \lambda) & C_4(1, \lambda) \end{vmatrix}.$$

Analogously to (4.5), we obtain the relation

$$\Delta^+(\lambda) = -D^+(\rho) \det(\mathcal{A}(\rho)), \quad (5.4)$$

where

$$D^+(\rho) = \begin{vmatrix} y_1(0, \lambda) & y_2(0, \lambda) & y_3(0, \lambda) & y_4(0, \lambda) \\ y_1'(0, \lambda) & y_2'(0, \lambda) & y_3'(0, \lambda) & y_4'(0, \lambda) \\ y_1^{[3]}(0, \lambda) & y_2^{[3]}(0, \lambda) & y_3^{[3]}(0, \lambda) & y_4^{[3]}(0, \lambda) \\ y_1(1, \lambda) & y_2(1, \lambda) & y_3(1, \lambda) & y_4(1, \lambda) \end{vmatrix}.$$

Similarly to (4.7), we derive

$$\begin{aligned} D^+(\rho) &= \rho^4 \exp(\rho\omega_4) d^+(\rho), & (5.5) \\ d^+(\rho) &= r_1^+(\rho) - r_2^+(\rho) \exp(\rho(\omega_3 - \omega_4)) + \varepsilon(\rho), \\ r_1^+(\rho) &= 4i - \frac{i\theta}{\rho} - \frac{2i\sigma}{\rho^2} - \frac{i(t_0 - t_1)}{2\rho^2} + \frac{i\theta^2}{8\rho^2}, \\ r_2^+(\rho) &= 4i + \frac{\theta}{\rho} + \frac{2i\sigma}{\rho^2} + \frac{i(t_0 - t_1)}{2\rho^2} - \frac{i\theta^2}{8\rho^2}. \end{aligned}$$

Substituting (4.5), (4.7), (5.4), and (5.5) into (5.1) for  $k = 3$ , we obtain

$$\beta_{n,3} = 4\lambda_{n,3} \frac{d^+(\rho_n)}{\frac{d}{d\rho} d(\rho_n)}, \quad (5.6)$$

where  $\lambda_{n,3} = \rho_n^4$  and  $\rho_n$  satisfies (4.8). Calculations show that

$$\begin{aligned} \frac{d^+(\rho)}{\frac{d}{d\rho} d(\rho)} &= \frac{r_1^+(\rho)r_2(\rho) - r_2^+(\rho)r_1(\rho)}{\frac{d}{d\rho} r_1(\rho)r_2(\rho) - \frac{d}{d\rho} r_2(\rho)r_1(\rho) - (\omega_3 - \omega_4)r_1(\rho)r_2(\rho)} + \varepsilon(\rho) \\ &= - \left( 1 + \frac{i(t_0 + \theta)}{4\rho^2} + \varepsilon(\rho) \right). \end{aligned} \quad (5.7)$$

Combining (4.8), (5.6), and (5.7), we arrive at (5.3) for  $\beta_{n,3}$ .  $\square$

## References

1. Aliyev Z.S., Kerimov N.B., Mehrabov V.A. Convergence of eigenfunction expansions for a boundary value problem with spectral parameter in the boundary conditions. I. *Differential Equations*, 2020, vol. 56, no. 2, pp. 143–157. <https://doi.org/10.1134/S0012266120020019>
2. Badanin A., Korotyaev E. Sharp eigenvalue asymptotics for fourth order operators on the circle. *Journal of Mathematical Analysis and Applications*, 2014, vol. 417, no. 2, pp. 804–818. <https://doi.org/10.1016/j.jmaa.2014.03.069>
3. Badanin A., Korotyaev E. Inverse problems and sharp eigenvalue asymptotics for Euler-Bernoulli operators. *Inverse Problems*, 2015, vol. 31, no. 5, 055004. <https://doi.org/10.1088/0266-5611/31/5/055004>
4. Barcion V. On the uniqueness of inverse eigenvalue problems. *Geophysical Journal International*, 1974, vol. 38, no. 2, pp. 287–298. <https://doi.org/10.1111/j.1365-246X.1974.tb04121.x>

5. Bondarenko N.P. Spectral data asymptotics for the higher-order differential operators with distribution coefficients. *Journal of Mathematical Sciences*, 2022, vol. 266, no. 5, pp. 794–815. <https://doi.org/10.1007/s10958-022-06118-x>
6. Bondarenko N.P. Linear differential operators with distribution coefficients of various singularity orders. *Mathematical Methods in the Applied Sciences*, 2023, vol. 46, no. 6, pp. 6639–6659. <https://doi.org/10.1002/mma.8929>
7. Bondarenko N.P. Regularization and inverse spectral problems for differential operators with distribution coefficients. *Mathematics*, 2023, vol. 11, no. 16, Article ID 3455. <https://doi.org/10.3390/math11163455>
8. Bondarenko N.P. Local solvability and stability of an inverse spectral problem for higher-order differential operators. *Mathematics*, 2023, vol. 11, no. 18, Art. ID 3818. <https://doi.org/10.3390/math11183818>
9. Bondarenko N.P. Necessary and sufficient conditions for solvability of an inverse problem for higher-order differential operators. *Mathematics*, 2024, vol. 12, no. 1, Article ID 61. <https://doi.org/10.3390/math12010061>
10. Gladwell G.M.L. Inverse Problems in Vibration. 2nd Ed. *Solid Mechanics and Its Applications*, vol. 119, Springer, Dordrecht, 2005.
11. Leibenzon Z.L. Algebraic-differential transformations of linear differential operators of arbitrary order and their spectral properties applicable to the inverse problem. *Mathematics of the USSR-Sbornik*, 1972, vol. 18, no. 3, pp. 425–471. <https://doi.org/10.1070/sm1972v018n03abeh001832>
12. McLaughlin J.R. An inverse eigenvalue problem of order four — an infinite case. *SIAM Journal of Mathematical Analysis*, 1978, vol. 9, no. 3, pp. 395–413. <https://doi.org/10.1137/0509026>
13. Mikhailets V., Molyboga V. Uniform estimates for the semi-periodic eigenvalues of the singular differential operators. *Methods of Functional Analysis and Topology*, 2004, vol. 10, no. 4, pp. 30–57.
14. Mikhailets V.A., Molyboga V.M. On the spectrum of singular perturbations of operators on the circle. *Mathematical Notes*, 2012, vol. 91, no. 4, pp. 588–591. <https://doi.org/10.1134/S0001434612030352>
15. Mirzoev K.A., Shkalikov A.A. Differential operators of even order with distribution coefficients. *Mathematical Notes*, 2016, vol. 99, no. 5, pp. 779–784. <https://doi.org/10.1134/S0001434616050163>
16. Möller M., Zinsou B. Asymptotics of the eigenvalues of self-adjoint fourth order differential operators with separated eigenvalue parameter dependent boundary conditions. *Rocky Mountain Journal of Mathematics*, 2017, vol. 47, no. 6, pp. 2013–2042. <https://doi.org/10.1216/RMJ-2017-47-6-2013>
17. Naimark M.A. *Linear Differential Operators, Part 1: Elementary Theory of Linear Differential Operators*. Ungar, New York, 1968.
18. Peletier L.A., Troy W.C. Spatial Patterns: Higher Order Models in Physics and Mechanics. Progress in Nonlinear Differential Equations and Their Applications, 2001, vol. 45, Birkhäuser, Boston. <https://doi.org/10.1007/978-1-4612-0135-9>
19. Polyakov D.M. Sharp eigenvalue asymptotics of fourth-order differential operators. *Asymptotic Analysis*, 2022, vol. 130, no. 3–4, pp. 477–503. <https://doi.org/10.3233/ASY-221760>
20. Polyakov D.M. On the spectral properties of a fourth-order self-adjoint operator. *Differential Equations*, 2023, vol. 59, no. 2, pp. 168–173. <https://doi.org/10.1134/S0012266123020027>
21. Polyakov D.M. Spectral asymptotics and a trace formula for a fourth-order differential operator corresponding to thin film equation. *Monatshefte für Mathematik*, 2023, vol. 202, no. 1, pp. 171–212. <https://doi.org/10.1007/s00605-022-01808-9>



22. Tamarkin J. Some general problems of the theory of ordinary linear differential equations and expansion of an arbitrary function in series of fundamental functions. *Mathematische Zeitschrift*, 1928, vol. 27, pp. 1–54. <https://doi.org/10.1007/BF01171084>
23. Vladimirov A.A. On one approach to definition of singular differential operators. *Cornell University, ArXiv*, 2017. <https://doi.org/10.48550/arXiv.1701.08017>
24. Yurko V.A. Method of Spectral Mappings in the Inverse Problem Theory. *Inverse and Ill-Posed Problems Series*, 2002, vol. 31, VNU Science, Utrecht. <https://doi.org/10.1515/9783110940961>
25. Yurko V.A. Asymptotics of solutions of differential equations with a spectral parameter. *Cornell University, ArXiv*, 2022. <https://doi.org/10.48550/arXiv.2204.07505>

### Об авторах

**Бондаренко Наталья Павловна,**

д-р физ.-мат. наук, доц.,  
Саратовский государственный  
университет, Саратов, 410012,  
Российская Федерация,  
[bondarenkonp@sgu.ru](mailto:bondarenkonp@sgu.ru),  
<https://orcid.org/0000-0003-2513-1472>

### About the authors

**Natalia P. Bondarenko,** Dr. Sci.  
(Phys.–Math.), Assoc. Prof., Saratov  
State University, Saratov, 410012,  
Russian Federation,  
[bondarenkonp@sgu.ru](mailto:bondarenkonp@sgu.ru),  
<https://orcid.org/0000-0003-2513-1472>

*Поступила в редакцию / Received 10.11.2023*

*Поступила после рецензирования / Revised 26.11.2023*

*Принята к публикации / Accepted 11.12.2023*