K. N. Ilinski, A. S. Stepanenko, q-Deformed superspace and q-extended supersymmetric Hamiltonian with arbitrary superpotential,
Зап. научн. сем. ПОМИ, 1996, том 235, 260–272

K. N. Ilinski, A. S. Stepanenko

q–DEFORMED SUPERSPACE AND q–EXTENDED SUPERSYMMETRIC HAMILTONIAN WITH ARBITRARY SUPERPOTENTIAL

§1. INTRODUCTION

Recently deformations of standard mathematical objects have attracted a lot of attention ([1,2] and refs. in [2]). In particular, in [3] the quantum group representation theory and q-deformed spaces have been studied and in [4] the procedure of supergroup quantization has been presented.

The attempts to construct the description of dynamics on q-deformed spaces were undertaken in [5, 6, 7]. In [8] q-deformed quantum spaces were considered as a gradedcommutative algebra. On this basis the classical and quantum dynamics on q-deformed spaces were proposed. The particles appeared after the quantization satisfy the q-deformed commutation relations:

$$a_i a_i^+ \mp a_i^+ a_i = 1 \qquad a_i a_j^+ \mp q a_j^+ a_i = 0$$

$$a_i^+ a_j^+ \mp q^{-1} a_j^+ a_i^+ = 0 \qquad a_i a_j \mp q^{-1} a_j a_i = 0$$

$$q^m = 1 \quad , \quad m \in N \quad , \quad 1 \leq i \leq j \leq n$$

$$a_i^{+2} = a_i^2 = 0 \quad \text{for} q\text{-deformed fermions},$$

where the upper sign corresponds to the quantization on q-deformed usual spaces and lower one corresponds to the quantization on q-deformed superspaces. The particles can be naturally called q-bosons and q-fermions. It is interesting to consider the generalization of supersymmetric theory for the case when q-fermions are included into the theory instead of fermions.

As we can see later such supersymmetric theory will differ from the deformed supersymmetry (q-supersymmetry or q-SUSY) by V. P .

Permanent address: Theoretical Department, St.-Petersburg Nuclear Physics Institute, Gatchina, St.-Petersburg, 188350, Russian Federation.

Spiridonov [9] where the following relations for the supercharges Q^+, Q^- were used:

$$qQ^+Q^- + q^{-1}Q^-Q^+ = H \quad {Q^+}^2 = {Q^-}^2 = 0$$
$$qHQ^- - q^{-1}Q^-H = qQ^+H - q^{-1}HQ^+ = 0 .$$

In our approach it is possible to discover the q-deformation only for the pair of different q-fermions and that is why this theory is q-extended supersymmetry (q-ESUSY). It is possible to coincide such theory in the framework of colour supersymmetry [10] (we are grateful N. Borisov for this remark).

In this paper we consider one of the possible generalization of SUSY. For this purpose we formulate a q-superspace formalism and construct the q-supertransformation group on this q-superspace. After this we start from a standard action for a scalar d-dimensional real superfield and build q-extended supersymmetric Lagrangian for component variables of the q-superfield. After the quantization procedure corresponding conserved quantities become q-supercharges. In difference from [11] where only general constructions of q-extended supersymmetry and simplest examples were considered in this paper we give exact expressions for q-supercharges and q-extended supersymmetric Hamiltonian with arbitrary superpotential.

§2. q-EXTENDED SUPERSYMMETRY AND q-DEFORMED SUPERSPACE FORMALISM

As in usual case [13, 14, 15] the most natural way to build supersymmetric Lagrangians is by employing the superspace formalism. For the sake of simplicity we will consider $(0+1)$-dimensional case, but all constructions can be generalized for multidimensional one.

For the $(0+1)$-dimensional case superspace reduces to the supertime. In the standard way besides time t complex anticommuting variables $\theta_i, \bar{\theta}_i, i = 1, 2$ are introduced. Instead of that we will consider the variables $\theta_i, \bar{\theta}_i \in HQ_m^{0|2}, i = 1, 2$, i.e.,

$$\theta_i\bar{\theta}_i = -\bar{\theta}_i\theta_i \qquad {\theta_i}^2 = \bar{\theta}_i^2 = 0 \qquad q^m = 1 \quad , \quad m \in N$$
$$\theta_1\theta_2 = -q^{-1}\theta_2\theta_1 \qquad \bar{\theta}_1\theta_2 = -q\theta_2\bar{\theta}_1$$
$$\bar{\theta}_1\bar{\theta}_2 = -q^{-1}\bar{\theta}_2\bar{\theta}_1 \qquad \theta_1\bar{\theta}_2 = -q\bar{\theta}_2\theta_1$$

It was shown in [8] that the algebra $HQ_m^{0|2}$ can be described as a graded-commutative algebra. For that we began from the algebra

$C(\theta_1, \bar{\theta}_1, \theta_2, \bar{\theta}_2)$ which is freely generated by $\theta_1, \bar{\theta}_1, \theta_2, \bar{\theta}_2$ and used a special graduation map. In this case the graduation map $\varphi : HQ_m^{0|2} \to Z_2 \oplus Z_m \oplus Z_m$ was defined by

$$\varphi(\theta_1^{k_1} \dots \bar{\theta}_2^{k_4}) = k_1\varphi(\theta_1) + \dots + k_4\varphi(\bar{\theta}_2)$$

$$\varphi(\theta_1) = (1,1,0)\ ; \quad \varphi(\theta_2) = (1,0,1)\ ; \quad \varphi(\bar{\theta}_i) = -\varphi(\theta_i)$$

and the graded-commutator for the homogeneous elements A, B had the form:

$$[A,B]_q^{0|2} = AB - (-1)^{\vartheta_2(\varphi(A),\varphi(B))} q^{-\vartheta_m(\varphi(A),\varphi(B))} BA$$

$$\vartheta_2(\varphi(A),\varphi(B)) = \varphi(A)|_{Z_2}\, \varphi(B)|_{Z_2}$$

and $\vartheta_m(\alpha,\beta)$ was defined by the following relations: $\vartheta_m : Z_2 \oplus Z_m \oplus Z_m \to Z$ being biadditive antisymmetric map,

$$\vartheta_m(\alpha,\beta) = \sum_{i,j} \alpha_i\beta_j\vartheta_m(\xi_i,\xi_j) \qquad \alpha = \sum_i \alpha_i\xi_i \qquad \beta = \sum_i \beta_i\xi_i$$

$$\vartheta_m(\xi_i,\xi_j) = 1 \quad i<j \qquad \xi_1 = (l,1,0) \qquad \xi_2 = (k,0,1) \qquad \forall k,l = 0,1$$

Our algebra $HQ_m^{0|2}$ can be described as a factor-algebra of $C(\theta_1, \bar{\theta}_1, \theta_2, \bar{\theta}_2)$ under the two-side ideal, generated by the elements $\{[C_1,C_2]_q^{0|2} \mid C_1, C_2 \in C(\theta_1 \dots \bar{\theta}_2)\}$ and in this sense $HQ_m^{0|2}$ is exactly the algebra with zero graded-commutators:

$$f,g \in HQ_m^{0|2} \qquad [f,g]_q^{0|2} = 0.$$

We note that q is a particular case of ϵ-factor [12] (and refs. therein).

A dynamic q-supervariable is a quantity depending on $t, \theta_1, \bar{\theta}_1, \theta_2, \bar{\theta}_2$. The coefficients of its expansion in $\theta_1, \bar{\theta}_1, \theta_2, \bar{\theta}_2$ have the commutation relations according to their graduations. In what follows we will use the following notations:

$$\theta = \begin{pmatrix}\theta_1\\ \theta_2\end{pmatrix} \qquad \bar{\theta} = (\bar{\theta}_1, \bar{\theta}_2) \qquad \psi = \begin{pmatrix}\psi_1\\ \psi_2\end{pmatrix} \qquad \bar{\psi} = (\bar{\psi}_1, \bar{\psi}_2)$$

$$\bar{\theta}\psi \equiv \bar{\theta}_1\psi_1 + \bar{\theta}_2\psi_2 \qquad \chi = \begin{pmatrix}\chi_1\\ \chi_2\end{pmatrix} \qquad \bar{\chi} = (\bar{\chi}_1, \bar{\chi}_2) \qquad \partial_t = \frac{\partial}{\partial t}$$

The supersymmetry group is implemented in terms of $t, \theta, \bar{\theta}$ transformations in the following way:

$$Q_{\bar{\eta}} \ : \quad t \to t + \frac{i}{2}\bar{\eta}\theta \qquad \bar{\theta} \to \bar{\theta} + \bar{\eta}$$

$$Q_\eta \ : \quad t \to t - \frac{i}{2}\bar{\theta}\eta \qquad \theta \to \theta + \eta$$

where $\eta = \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix}$, $\bar{\eta} = (\bar{\eta_1}, \bar{\eta_2})$ are transformation parameters, such as $\varphi(\eta_i) = \varphi(\theta_i)$.

The generators of the supersymmetry transformations can be written in a differential form:

$$Q_{\bar{\eta}} = \frac{\partial}{\partial\theta} + \frac{i}{2}\theta\partial_t \equiv \begin{pmatrix} Q^1_{\bar{\eta}} \\ Q^2_{\bar{\eta}} \end{pmatrix} \qquad Q_\eta = -\frac{\partial}{\partial\theta} - \frac{i}{2}\bar{\theta}\partial_t \equiv (Q^1_\eta, Q^2_\eta) \qquad (1)$$

Now we note that

$$[Q^\alpha_{\bar{\eta}}, Q^\alpha_\eta]^{0|2}_q = -i\partial_t \qquad \alpha = 1, 2$$

and

$$[Q^1_\eta, Q^2_\eta]^{0|2}_q = Q^1_\eta Q^2_\eta + q^{-1} Q^2_\eta Q^1_\eta = 0 \qquad Q^{\alpha 2}_{\bar{\eta}} = Q^{\alpha 2}_\eta = 0$$

$$[Q^1_{\bar{\eta}}, Q^2_{\bar{\eta}}]^{0|2}_q = [Q^\alpha_{\bar{\eta}}, Q^{\alpha'}_\eta]^{0|2}_q = 0 \qquad \alpha' \neq \alpha. \qquad (2)$$

Above relations give the connections between the supersymmetry generators and the Hamiltonian of the system. So, graded-commutators $[Q^\alpha_{\bar{\eta}}, Q^\alpha_\eta]^{0|2}_q, \alpha = 1, 2$ are the generators of time translations (up to a factor $-i$).

Representation of the supersymmetric group can be implemented, for example, by means of d-dimensional scalar real q-superfield $\phi = (\phi^1 \dots \phi^d)$. Its expansion in $\theta, \bar{\theta}$ looks like

$$\begin{aligned} \phi^k(t) = x^k(t) + \frac{1}{\sqrt{2}}\bar{\theta}\psi^k(t) + \frac{1}{\sqrt{2}}\bar{\psi}^k(t)\theta + \\ + \bar{L}^k(t)\theta_1\theta_2 + \bar{\theta}_2\bar{\theta}_1 L^k(t) + \mathrm{Tr}\,(\widehat{\theta}\mathbf{F}^k(t)) + \frac{1}{\sqrt{2}}\bar{\chi}^k(t)\tilde{\theta} + \\ + \frac{1}{\sqrt{2}}\tilde{\bar{\theta}}\chi^k(t) + H^k(t)\bar{\theta}_1\theta_1\bar{\theta}_2\theta_2\,, \qquad k = 1 \dots d \end{aligned} \qquad (3)$$

where

$$\widehat{\theta} = \begin{pmatrix} \bar{\theta}_1\theta_1 & \bar{\theta}_2\theta_1 \\ \bar{\theta}_1\theta_2 & \bar{\theta}_2\theta_2 \end{pmatrix}, \quad \mathbf{F} = \begin{pmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{pmatrix}, \quad \widehat{\theta} = \begin{pmatrix} \theta_1\bar{\theta}_2\theta_2 \\ \theta_2\bar{\theta}_1\theta_1 \end{pmatrix}.$$

Q-supersymmetry transformations are realized in component variables as follows:

for $\bar{\eta}Q_{\bar{\eta}}$

$$\delta_1 x^k = \frac{1}{\sqrt{2}}\bar{\eta}\psi^k \quad , \quad \delta_1 H^k = -\frac{i}{2\sqrt{2}}\bar{\eta}\dot{\chi}^k$$

$$\delta_1\bar{\psi}^k = \sqrt{2}\left\{\frac{i}{2}\bar{\eta}\dot{x}^k + (\bar{\eta}_1 F_{11}^k + \bar{\eta}_2 q^{-1} F_{21}^k \ , \ \bar{\eta}_2 F_{22}^k + \bar{\eta}_1 q F_{12}^k)\right\},$$

$$\delta_1\psi^k = \sqrt{2}\begin{pmatrix} -q\bar{\eta}_2 \\ \bar{\eta}_1 \end{pmatrix} L^k$$

$$\delta_1\bar{\chi}^k = \sqrt{2}\left\{\bar{\eta}H^k + \frac{i}{2}(\bar{\eta}_1\dot{F}_{22}^k - \bar{\eta}_2 q^{-1}\dot{F}_{21}^k \ , \ \bar{\eta}_2\dot{F}_{11}^k - \bar{\eta}_1 q\dot{F}_{12}^k)\right\},$$

$$\delta_1\chi^k = -\frac{i}{\sqrt{2}}\begin{pmatrix} -q\bar{\eta}_2 \\ \bar{\eta}_1 \end{pmatrix}\dot{L}^k$$

$$\delta_1 L^k = 0, \quad \delta_1\bar{L}^k = \frac{1}{\sqrt{2}}\left(\bar{\eta}_2(\frac{i}{2}\dot{\bar{\psi}}_1^k + \bar{\chi}_1^k) + (\frac{i}{2}\dot{\bar{\psi}}_2^k + \bar{\chi}_2^k)\bar{\eta}_1\right)$$

$$\delta_1\mathbf{F}^k = \delta_1\begin{pmatrix} F_{11}^k & F_{12}^k \\ F_{21}^k & F_{22}^k \end{pmatrix} =$$

$$= \frac{1}{\sqrt{2}}\begin{pmatrix} \bar{\eta}_1(-\frac{i}{2}\dot{\psi}_1^k) + \bar{\eta}_2\chi_2^k & -\bar{\eta}_2(\frac{i}{2}\dot{\psi}_1^k + \chi_1^k) \\ -\bar{\eta}_1(\frac{i}{2}\dot{\psi}_2^k + \chi_2^k) & \bar{\eta}_2(-\frac{i}{2}\dot{\psi}_2^k) + \bar{\eta}_1\chi_1^k \end{pmatrix} \tag{4}$$

and for $Q_\eta\eta$ it is possible to obtain transformation laws using (1) and (4).

The covariant derivatives D and $\bar{D}$ have the form

$$D = \frac{\partial}{\partial\bar{\theta}} - \frac{i}{2}\theta\partial_t \equiv \begin{pmatrix} D_1 \\ D_2 \end{pmatrix} \qquad \bar{D} = -\frac{\partial}{\partial\theta} + \frac{i}{2}\bar{\theta}\partial_t \equiv (\bar{D}_1, \bar{D}_2)$$

satisfying the usual conditions

$$D\delta_i\phi = \delta_i D\phi \ , \qquad \bar{D}\delta_i\phi = \delta_i\bar{D}\phi$$

where $\delta_i\phi, i = 1,2$ are the variances of the q-supervariables under a supersymmetric transformations.

The action for the q-extended supersymmetric quantum mechanics can be written in the following way

$$S = \int \mathrm{d}\bar{\theta}_1\mathrm{d}\theta_1\mathrm{d}\bar{\theta}_2 d\theta_2 \ \left(\bar{D}\phi^k D\phi^k - 2V(\phi)\right) = \int dt \ \mathcal{L}$$

where $V(\cdot)$ is an arbitrary function (superpotential). We imply the integration over $\theta_i, \bar{\theta}_i$ in the following sense

$$\int 1\mathrm{d}\theta_i = \int 1\mathrm{d}\bar{\theta}_i = \int \theta_j \mathrm{d}\bar{\theta}_i = \int \bar{\theta}_j \mathrm{d}\theta_i = 0 \qquad \forall i,j = 1,2$$

$$\int \theta_i \mathrm{d}\theta_k = \int \bar{\theta}_i \mathrm{d}\bar{\theta}_k = \delta_{ik}\ , \qquad \phi(\theta_i) = \phi(\mathrm{d}\theta_i)\ , \qquad \phi(\bar{\theta}_i) = \phi(\mathrm{d}\bar{\theta}_i)$$

As a result the Lagrangian takes the form

$$\begin{aligned}
\mathcal{L} = &-\frac{1}{2}\dot{x}^k\left(\dot{F}^k_{11}+\dot{F}^k_{22}\right) - 2H^k\left(F^k_{11}+F^k_{22}\right) + \frac{1}{4}\dot{\bar{\psi}}^k\dot{\psi}^k + \bar{\chi}^k\chi^k + \\
&+\frac{i}{4}\left(\dot{\bar{\psi}}^k\chi^k - \bar{\psi}^k\dot{\chi}^k + \dot{\bar{\chi}}^k\psi^k - \bar{\chi}^k\dot{\psi}^k\right) + 2i\bar{L}^k\dot{L}^k - 2\frac{\partial V}{\partial x^k}H^k + \\
&+2\frac{\partial^2 V}{\partial x^k \partial x^l}\left(\frac{1}{2}\bar{\psi}^l\chi^k + \frac{1}{2}\bar{\chi}^l\psi^k + F^k_{12}F^l_{21} - F^k_{11}F^l_{22} - \bar{L}^l L^k\right) - \\
&-\frac{\partial^3 V}{\partial x^l \partial x^m \partial x^n}\Big(F^l_{12}\bar{\psi}^m_1\psi^n_2 + F^l_{21}\bar{\psi}^m_2\psi^n_1 - F^l_{11}\bar{\psi}^m_2\psi^n_2 - F^l_{22}\bar{\psi}^m_1\psi^n_1 + \\
&+\bar{L}^l\psi^m_1\psi^n_2 + L^l\bar{\psi}^m_2\bar{\psi}^n_1\Big) - \frac{1}{2}\frac{\partial^4 V}{\partial x^l\partial x^m\partial x^n\partial x^k}\bar{\psi}^l_1\bar{\psi}^m_2\psi^n_2\psi^k_1 \qquad (6)
\end{aligned}$$

where full derivatives have been neglected. The auxiliary variables F^k_{ij}, χ^k, $\bar{\chi}^k$, H^k may be excluded in the usual way if matrix $\partial_{lk}V = K$ is invertible. Omitting spatial symbols to simplify the expressions we can put down the expressions for auxiliary fields

$$\begin{aligned}
\chi_i(t) &= (-2\psi_i(t)V''[x(t)] + \dot{\bar{\psi}}_i(t)i)/2 \\
\bar{\chi}_i(t) &= -(2\bar{\psi}_1(t)V''[x(t)] + \dot{\bar{\psi}}_i(t)i)/2 \\
F_{i,j\neq i}(t) &= K^{-1}V'''[x(t)]\bar{\psi}_j(t)\psi_i(t)/2 \\
F_{ii}(t) &= (-1)^i\frac{1}{4}((-1)^k\bar{\psi}_k(t)\psi_k(t)K^{-1}V'''[x(t)]) - \frac{1}{2}V'[x(t)] \\
H(t) &= \frac{1}{4}(\bar{\psi}_i(t)\psi_i(t)V'''[x(t)]) + \frac{1}{2}V'[x(t)]V''[x(t)] + \frac{1}{4}\ddot{x}(t).
\end{aligned}$$

After substitutions of auxiliary fields Lagrangian has the form:

$$\mathcal{L} = -\Big(8\bar{L}(t)L(t)V''[x(t)] - 8\bar{L}(t)\dot{L}(t)i + 4\bar{L}(t)\psi_1(t)\psi_2(t)V'''[x(t)] -$$
$$-4\bar{\psi}_1(t)\bar{\psi}_2(t)L(t)V'''[x(t)]q + 3\bar{\psi}_1(t)\bar{\psi}_2(t)\psi_1(t)\psi_2(t)K^{-1}\left(V'''[x(t)]\right)^2 q -$$
$$-2\bar{\psi}_1(t)\bar{\psi}_2(t)\psi_1(t)\psi_2(t)V^{(IV)}[x(t)]q + 2\bar{\psi}_i(t)\psi_i(t)V'[x(t)]V'''[x(t)] +$$
$$+4\bar{\psi}_i(t)\psi_i(t)\left(V''[x(t)]\right)^2 - 2\bar{\psi}_i(t)\dot{\psi}_i(t)V''[x(t)]i + 2\dot{\bar{\psi}}_i(t)\psi_i(t)V''[x(t)]i +$$
$$+2\left(V'[x(t)]\right)^2 V''[x(t)] - 2V''[x(t)]\left(\dot{x}(t)\right)^2\Big)/4.$$

Let us denote the action as follows

$$S = \int \mathrm{d}t\mathrm{d}\bar{\theta}_1\mathrm{d}\theta_1\mathrm{d}\bar{\theta}_2\mathrm{d}\theta_2\ \mathcal{L} = \int \mathrm{d}t\ \widehat{\mathcal{L}}\,.$$

We can find conserved Noether's currents directly for $\widehat{\mathcal{L}}$, then integrate over $\theta, \bar{\theta}$ and obtain conserved supercharges. These expressions contain not only "physical" fields but also auxiliary fields. After substitutions of constrains supercharges have the following form in terms of "physical" fields:

$$Q_{\eta_i}\eta_i = -\left(4\bar{L}(t)\psi_2(t)V''[x(t)]iq^{-1} - \bar{\psi}_1(t)\bar{\psi}_2(t)\psi_2(t)V'''[x(t)]i -\right.$$
$$\left. - 2\bar{\psi}_1(t)V'[x(t)]V''[x(t)]i + 2\bar{\psi}_1(t)V''[x(t)]\dot{x}(t)\right)\eta_1/2\sqrt{2}$$
$$Q_{\eta_2}\eta_2 = \left(4\bar{L}(t)\psi_1(t)V''[x(t)]i - \bar{\psi}_1(t)\bar{\psi}_2(t)\psi_1(t)V'''[x(t)]iq +\right.$$
$$\left. + 2\bar{\psi}_2(t)V'[x(t)]V''[x(t)]i - 2\bar{\psi}_2(t)V''[x(t)]\dot{x}(t)\right)\eta_2/2\sqrt{2}$$
$$Q_{\bar{\eta}_1}\bar{\eta}_1 = -\left(4\bar{\psi}_2(t)L(t)V''[x(t)]iq + \bar{\psi}_2(t)\psi_1(t)\psi_2(t)V'''[x(t)]iq -\right.$$
$$\left. - 2\psi_1(t)V'[x(t)]V''[x(t)]i - 2\psi_1(t)V''[x(t)]\dot{x}(t)\right)\bar{\eta}_1/2\sqrt{2}$$
$$Q_{\bar{\eta}_2}\bar{\eta}_2 = \left(4\bar{\psi}_1(t)L(t)V''[x(t)]i + \bar{\psi}_1(t)\psi_1(t)\psi_2(t)V'''[x(t)]i +\right.$$
$$\left. + 2\psi_2(t)V'[x(t)]V''[x(t)]i + 2\psi_2(t)V''[x(t)]\dot{x}(t)\right)\bar{\eta}_2/2\sqrt{2}.$$

§3. Classical dynamics and quantization procedure for the component variables

Here we construct classical and quantum dynamics in close analogy with [8]. It is important however that the new fields $L, \bar{L}$ arise which are not q-fields in the sense [8].

To sake of simplicity let us make the change of variables:

$$\bar{L},\ L \to \frac{\bar{L}}{\sqrt{2}},\ \frac{L}{\sqrt{2}}\ ,\quad \bar{\psi} \to \bar{\psi}K^{-1/2},\ \psi K^{-1/2}.$$

Then we have for Lagrangian of the system can be rewritten in the form:

$$\mathcal{L} = -\Big(4\bar{L}(t)L(t)V''[x(t)] - 2\bar{L}(t)\dot{L}(t)i + 2\dot{\bar{L}}(t)L(t)i +$$
$$+2\sqrt{2}\bar{L}(t)\psi_1(t)\psi_2(t)K^{-1}V'''[x(t)] -$$
$$-2\sqrt{2}\bar{\psi}_1(t)\bar{\psi}_2(t)L(t)K^{-1}V'''[x(t)]q +$$
$$+3\bar{\psi}_1(t)\bar{\psi}_2(t)\psi_1(t)\psi_2(t)K^{-3}\left(V'''[x(t)]\right)^2 q +$$
$$+2\bar{\psi}_i(t)\psi_i(t)K^{-1}V'[x(t)]V'''[x(t)] -$$
$$-2\bar{\psi}_1(t)\bar{\psi}_2(t)\psi_1(t)\psi_2(t)K^{-2}V^{(IV)}[x(t)]q +$$
$$+4\bar{\psi}_i(t)\psi_i(t)K - 2\bar{\psi}_i(t)\dot{\psi}_i(t)i +$$
$$+2\dot{\bar{\psi}}_i(t)\psi_i(t)i + 2\left(V'[x(t)]\right)^2 K - 2K\dot{x}^2(t)\Big)/4.$$

The principle of extreme action for trajectories is

$$\delta S = 0$$

$$S[x,p,\psi,\bar{\psi},L,\bar{L}] = \int \mathrm{d}t \left\{\frac{i}{2}\sum_{j=1}^{2}[\bar{\psi}_j(t)\dot{\psi}_j(t) - \dot{\bar{\psi}}_j(t)\psi_j(t)] + \right.$$
$$\left. + \frac{i}{2}[\bar{L}(t)\dot{L}(t) - \dot{\bar{L}}(t)L(t)] + p(t)\dot{x}(t) - \mathcal{H}[x,p,\psi,\bar{\psi},L,\bar{L}]\right\} \equiv \int \mathrm{d}t\ \mathcal{L}$$

where

$$\dot{\psi}_j^{\,k} = \frac{d\psi_j^k}{dt}, \quad \dot{\bar{\psi}}_j^{\,k} = \frac{d\bar{\psi}_j^{\,k}}{dt}, \quad \dot{L}^k = \frac{dL^k}{dt}, \quad \dot{\bar{L}}^k = \frac{d\bar{L}^k}{dt}, \quad \varphi(\mathcal{H}) = 0.$$

Derivatives are calculated as in the Grassmann algebra case.

The equations for extreme trajectories have the form:

$$\frac{\partial\mathcal{L}}{\partial\psi_j} = \frac{\mathrm{d}}{\mathrm{d}t}\frac{\partial\mathcal{L}}{\partial\dot{\psi}_j}, \quad \frac{\partial\mathcal{L}}{\partial\bar{\psi}_j} = \frac{\mathrm{d}}{\mathrm{d}t}\frac{\partial\mathcal{L}}{\partial\dot{\bar{\psi}}_j}, \quad \frac{\partial\mathcal{L}}{\partial L} = \frac{\mathrm{d}}{\mathrm{d}t}\frac{\partial\mathcal{L}}{\partial\dot{L}},$$

$$\frac{\partial\mathcal{L}}{\partial\bar{L}} = \frac{\mathrm{d}}{\mathrm{d}t}\frac{\partial\mathcal{L}}{\partial\dot{\bar{L}}}, \quad \frac{\partial\mathcal{L}}{\partial x} = \frac{\mathrm{d}}{\mathrm{d}t}\frac{\partial\mathcal{L}}{\partial\dot{x}}, \quad \frac{\partial\mathcal{L}}{\partial p} = \frac{\mathrm{d}}{\mathrm{d}t}\frac{\partial\mathcal{L}}{\partial\dot{p}}.$$

Here $\frac{\partial}{\partial L^k}, \frac{\partial}{\partial\bar{L}^k}, \frac{\partial}{\partial\psi_j{}^k}, \frac{\partial}{\partial\bar{\psi}_j{}^k}$ are defined as left derivatives in close analogy with the Grassmann case. Note that they are differentiations of graduated algebra of functions, i.e.,

$$\frac{\partial}{\partial\psi_j{}^k}fg = \left(\frac{\partial}{\partial\psi_j{}^k}f\right)g + (-1)^{-\theta_2(\varphi(\psi_j^k),\varphi(f))}q^{-\theta_m(\varphi(\bar{\psi}_j{}^k),\varphi(f))}f\left(\frac{\partial}{\partial\psi_j{}^k}g\right)$$

for homogeneous elements f, g and similar for $\frac{\partial}{\partial \bar{L}}$, $\frac{\partial}{\partial L}$, $\frac{\partial}{\partial \bar{\psi}}$.

Putting $p = \dot{x}K$ we can rewrite the motion equations in Hamiltonian form:

$$\dot{\chi}_k = \left(\Omega^{-1}\right)_{kj} \frac{\partial \mathcal{H}}{\partial \chi_j}$$

where

$$\begin{pmatrix} \psi_1 \\ \bar{\psi}_1 \\ \psi_2 \\ \bar{\psi}_2 \\ L \\ \bar{L} \\ x \\ p \end{pmatrix}, \quad \Omega \equiv \begin{pmatrix} \Omega_1 & 0 & 0 & 0 \\ 0 & \Omega_2 & 0 & 0 \\ 0 & 0 & \Omega_3 & 0 \\ 0 & 0 & 0 & \Omega_4 \end{pmatrix},$$

$$\Omega_{1,2} \equiv \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad \Omega_3 \equiv \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \Omega_4 \equiv \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Hence, the inverse matrix Ω^{-1} is

$$\Omega^{-1} = \begin{pmatrix} \Omega_1^{-1} & 0 & 0 & 0 \\ 0 & \Omega_2^{-1} & 0 & 0 \\ 0 & 0 & \Omega_3^{-1} & 0 \\ 0 & 0 & 0 & \Omega_4^{-1} \end{pmatrix},$$

$$\Omega_{1,2}^{-1} = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}, \quad \Omega_3^{-1} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \Omega_4^{-1} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

It is possible to introduce the Poisson structure in close analogy with the Grassmann case:

$$\{f, g\}_{\mathrm{cl}} = -\sum_{s,s'} \left(\Omega^{-1}\right)_{s,s'} \frac{\partial}{\partial \chi'_{s'}} \frac{\partial}{\partial \chi_s} f(\chi) g(\chi') \Big|_{\chi=\chi'}.$$

Such defined Poisson brackets satisfy the following relations for homogeneous elements f, g, h:

$$\{f, g\}_{\mathrm{cl}} = (-1)^{\theta_2(\varphi(f),\varphi(g))} q^{\theta_m(\varphi(f),\varphi(g))} \{g, f\}_{\mathrm{cl}}$$

$$\{fg, h\}_{\mathrm{cl}} = f\{g, h\}_{\mathrm{cl}} + (-1)^{\theta_2(\varphi(g),\varphi(h))} q^{\theta_m(\varphi(g),\varphi(h))} \{f, h\} g$$

$$\{f, \{g, h\}_{\mathrm{cl}}\}_{\mathrm{cl}} +$$

$$+(-1)^{\theta_2(\varphi(g),\varphi(h))+\theta_2(\varphi(f),\varphi(h))} q^{\theta_m(\varphi(g),\varphi(h))+\theta_m(\varphi(f),\varphi(h))} \{h, \{f, g\}_{\mathrm{cl}}\}_{\mathrm{cl}} +$$

$$+(-1)^{\theta_2(\varphi(f),\varphi(g))+\theta_2(\varphi(f),\varphi(h))} q^{\theta_m(\varphi(f),\varphi(g))+\theta_m(\varphi(f),\varphi(h))} \{g, \{h, f\}_{\mathrm{cl}}\}_{\mathrm{cl}} = 0 \tag{7}$$

which correspond to generalization of antisymmetry, differentiation property and Jacobi identity for ordinary Poisson brackets.

The equations of motion can be written by means of these Poisson brackets as

$$\dot{\chi}_j = \{H, \chi_j\}_{\rm cl}.$$

It should be noted that classical dynamics complies with the graduation structure (i.e., evolution does not change graduation of χ) due to $\varphi(H) = 0$.

The commutation relations for canonical variables related to $\{\ \}_{\rm cl}$ have the form:

$$\{\psi_i, \bar{\psi}_i\}_{\rm cl} = i\ ,\quad \{L, \bar{L}\}_{\rm cl} = i\ ,\quad \{x, p\}_{\rm cl} = -1 \tag{8}$$

and others are zeroth.

To canonically quantize our theory one has

1 to construct the operators of canonical variables in Hilbert space, which satisfy the commutation relations (8)
2 to define the quantum Poisson brackets $\{\ \}_\hbar$ which have the same algebraic properties as the classical ones.

In this case item 2 leads to following definition of quantum Poisson brackets:

$$\{A, B\}_\hbar = \frac{i}{\hbar}[A, B]_q^{0|2}.$$

The commutation relations for the field operators follow from the above:

$$(L^+)^+ = L\ ,\quad (\psi_i^+)^+ = \psi_i\ ,\quad q^* = q^{-1}\ ,$$

$$(\psi_i^+)^2 = (\psi_i)^2 = 0\ ,\quad \psi_1\psi_2 + q^{-1}\psi_2\psi_1 = 0\ ,\quad \psi_1\psi_2^+ + q\psi_2^+\psi_1 = 0\ ,$$

$$L\psi_1 - q\psi_1 L = 0\ ,\quad L\psi_2 - q^{-1}\psi_2 L = 0\ ,$$

$$L\psi_2^+ - q\psi_2^+ L = 0\ ,\quad L\psi_1^+ - q^{-1}\psi_1^+ L = 0\ ,$$

$$LL^+ - L^+L = h\ ,\quad \psi_i\psi_i^+ + \psi_i^+\psi_i = h\ ,\quad [x, p] = ih \tag{9}$$

and relations which can be obtained by Hermitian conjugation.

After quantization q-supercharges have the form:

$$Q_1 = -(2\sqrt{2}\bar{L}\psi_2 K^{1/2} iq^{-1} - \bar{\psi}_1\bar{\psi}_2\psi_2 K^{-3/2}V'''i + \bar{\psi}_1 K^{-3/2}V'''hi +$$
$$+ 2\bar{\psi}_1 K^{-1/2}p - 2\bar{\psi}_1 K^{1/2}V'i)/2\sqrt{2}$$
$$Q_2 = (2\sqrt{2}\bar{L}\psi_1 K^{1/2}i - \bar{\psi}_1\bar{\psi}_2\psi_1 K^{-3/2}V'''iq - \bar{\psi}_2 K^{-3/2}V'''hi -$$
$$- 2\bar{\psi}_2 K^{-1/2}p + 2\bar{\psi}_2 K^{1/2}V'i)/2\sqrt{2}$$
$$Q_1^+ = (2\sqrt{2}\bar{\psi}_2 L K^{1/2}iq + \bar{\psi}_2\psi_1\psi_2 K^{-3/2}V'''iq -$$
$$- 2\psi_1 K^{-1/2}p - 2\psi_1 K^{1/2}V'i)/2\sqrt{2}$$
$$Q_2^+ = -(2\sqrt{2}\bar{\psi}_1 L K^{1/2}i + \bar{\psi}_1\psi_1\psi_2 K^{-3/2}V'''i +$$
$$+ 2\psi_2 K^{-1/2}p + 2\psi_2 K^{1/2}V'i)/2\sqrt{2}.$$

It is possible to check that such q-supercharges obey the q-extended supersymmetry algebra

$$\{Q_i, Q_i^+\} = H\ ,\quad (Q_i)^2 = (Q_i^+)^2 = 0\ ,\quad (Q_i^+)^+ = Q_i\ ,$$
$$Q_1Q_2 = -q^{-1}Q_2Q_1\ ,\quad Q_1Q_2^+ = -qQ_2^+Q_1\ ,\quad i = 1,2$$

with following quantum q- extended supersymmetric Hamiltonian

$$\mathcal{H} = h\Big(8\bar{L}LK + 4\sqrt{2}\bar{L}\psi_1\psi_2 K^{-1}V''' - 4\sqrt{2}\bar{\psi}_1\bar{\psi}_2 L K^{-1}V'''q +$$
$$+6\bar{\psi}_1\bar{\psi}_2\psi_1\psi_2 K^{-3}(V''')^2 q - 4\bar{\psi}_1\bar{\psi}_2\psi_1\psi_2 K^{-2}V^{(IV)}q +$$
$$+3\bar{\psi}_1\psi_1 K^{-3}(V''')^2 h - 2\bar{\psi}_1\psi_1 K^{-2}V^{(IV)}h + 4\bar{\psi}_1\psi_1 K^{-1}V'V''' +$$
$$+8\bar{\psi}_1\psi_1 K + 3\bar{\psi}_2\psi_2 K^{-3}(V''')^2 h - 2\bar{\psi}_2\psi_2 K^{-2}V^{(IV)}h +$$
$$+4\bar{\psi}_2\psi_2 K^{-1}V'V''' + 8\bar{\psi}_2\psi_2 K - 3K^{-3}(V''')^2 h^2 +$$
$$+4K^{-2}V'''phi + 2K^{-2}V^{(IV)}h^2 - 4K^{-1}V'V'''h +$$
$$+4K^{-1}p^2 + 4K(V')^2 - 4Kh\Big)/8.$$

Let us introduce the creation (annihilation) operators a_i^+, a_i, l^+, l which correspond to the fields $\psi, \bar{\psi}, L, \bar{L}$:

$$\psi_i^k = \sqrt{h}a_i^k \qquad L^k = \sqrt{h}l^k$$
$$\psi_i^{k+} = \sqrt{h}a_i^{k+} \qquad L^{k+} = \sqrt{h}l^{k+}$$

It is easy to obtain their commutation relations from (9).

Now we would like to discuss creation and annihilation operator representations. Let us consider the relations:

$$a_1^k = b_1^k\ , \qquad (a_1^{k+})^+ = a_1^k$$

$$a_2^k = e^{i\varphi\sum_{i=1}^d (b_1^i)^+ b_1^i} b_2^k\ , \qquad (a_2^{k+})^+ = a_2^k\ , \qquad q = e^{-i\varphi} \tag{10}$$

$$l^k = e^{i\varphi\sum_{i=1}^d (b_1^{i+} b_1^i - b_2^{i+} b_2^i)} d^k\ , \qquad (l^{k+})^+ = l^k$$

where b_i^{k+}, b_i^k, d^{k+}, d^k are creation (annihilation) operators of bosons and fermions respectively with usual commutation relations:

$$[b_i^k, d^l] = [b_i^{k+}, d^l] = [b_i^k, d^{l+}] = [b_i^{k+}, d^{l+}] = 0$$
$$\{b_i^k, b_j^{l+}\} = \delta_{ij} K_{kl}^{-1}\ , \qquad [d^k, d^{l+}] = M_{kl}^{-1}$$
$$\{b_i^k, b_j^l\} = \{b_i^{k+}, b_j^{l+}\} = 0\ , \qquad [d^k, d^l] = [d^{k+}, d^{l+}] = 0.$$

Using matrix representation for fermion operators we can put down matrix representation for q-fermion operators and operators l^k, l^{k+} can be implemented by means of matrix differential operators. Hence q-supercharges and q-extended supersymmetric Hamiltonian can be represented by means of matrix differential operators as it was done for usual supercharges and supersymmetric Hamiltonian in [14].

§4. Conclusion

In this paper we have studied the generalization of the supersymmetry, which was generated by changing fermions by q-fermions. The latters can be obtained after quantization of the fields, which take values in q-deformed spaces.

For this purpose we formulated the q-superspace formalism and constructed the q-supertransformation group. On this basis q-extended supersymmetric Lagrangian was built and corresponding q-supercharges were put down. We considered the rule for the quantization such systems and fulfilled this procedure for our case. We note that all q-deformed objects (q-supercharges and q-extended supersymmetric Hamiltonian) contain arbitrary functional parameter – superpotential V. It is especially interesting for investigations of topological properties of q-extended supersymmetric Hamiltonians.

References

1. L. D. Faddeev, N. Y. Reshetikhin, and L. A. Takhtajan, *Quantization of Lie groups and Lie algebras.* Preprint LOMI, 1987.
2. S. Majid, — Int. J. of Mod. Phys. A, **v.5, No. 1,** (1990), 1.
3. Yu. I. Manin, *Quantum groups and non-commutative geometry.* Preptint Montreal University, CRM-1561 1988.
4. Yu. I. Manin, — Comm. in Math. Phys. **123, No. 1** (1990), 163.
5. J. Wess, B. Zumino, Preprint CERN-TH-5697/90.
6. U. Carow-Watamura, M. Schlieker, and S. Watamura, Preprint KA-THEP-1990-15.
7. I. Ya. Aref'eva, I. V. Volovich, *Quantum group particles and Non-Archimedean geometry.* Preprint CERN-TH-6137/91.
8. N. V. Borisov, K. N. Ilinski, and V. M. Uzdin, — Phys. Lett. **A 169** (1992), 427.

no 9 V. P. Spiridonov, — Mod. Phys. Lett. **A 7** (1992), 1241.

10. V. Rittenberg, D. Wyler, Nucl. Phys..
11. K. N. Ilinski, V. M. Uzdin, *Quantum superspace, q-extended supersymmetry and parasupersymmetric quantum mecanics..* — submitted in Mod. Phys. Lett..
12. W. Marcinek, — J. Math. Phys. **33(5)** (1992), 1631.
13. V. P. Akulov, A. I. Pashkov, — Sov. J. Theor. Math. Phys. **56** (1983), 344.
14. A. A. Andrianov, N. V. Borisov, M. V. Ioffe, and M. I. Eides, — Teor. Mat. Fiz. **61, No. 1** (1984), 17.
15. A. V. Smilga, — Nucl. Phys. **B 249** (1985), 413.

St.-Petersburg
Steklov Mathematical Institute

Поступило 12 мая 1993г.

Service de Physique Théorique
CE-Saclay, F-91191
Gif-sur-Yvette Cedex, France